arXiv:hep-ph/0112159v2 13 Dec 2002

Brane-World Multigravity

Antonios Papazoglou
Keble College

Theoretical Physics
Department of Physics
University of Oxford

Thesis submitted for the degree of Doctor of Philosophy
in the University of Oxford

· September 2001 ·

Brane-World Multigravity

Antonios Papazoglou
Keble College

Theoretical Physics
Department of Physics
University of Oxford

Abstract

In this thesis, we discuss the idea of multigravity, namely the possibility that a significant component of gravity that we feel at intermediate distances ($1\text{mm} < r < 10^{26}\text{cm}$) is due to massive but ultralight gravitons. We demonstrate how this phenomenon can be realized in brane-world models in a spacetime with more than four dimensions and discuss how modifications of gravity at cosmological scales emerge as a consequence. Firstly, we consider five dimensional multigravity models with flat branes. We see how the existence of freely moving negative tension branes gives rise to ultralight graviton Kaluza-Klein states. Secondly, we study the moduli corresponding to the position of the branes and the size of the extra dimension, the radions and the dilaton respectively. We show that the radions corresponding to negative tension branes have wrong sign kinetic term. We also derive a stabilization condition for the dilaton in a brane model with general bulk scalar field dynamics. Thirdly, we show how we can reproduce the five dimensional multigravity models with Anti-de Sitter branes of positive tension only. Moreover, we demonstrate how the van Dam-Veltman-Zakharov discontinuity can be circumvented at tree level in Anti-de Sitter space. Finally, we discuss how the above multigravity models can be replicated in six dimensions with flat positive tension branes. This last possibility is the first theoretically and phenomenologically viable realization of the multigravity scenario.

Thesis submitted for the Degree of Doctor of Philosophy
in the University of Oxford

· September 2001 ·

To my parents, Eleni and Yiannis.

"Leave the beaten track behind occasionally and dive into the woods. Every time you do you will be certain to find something that you have never seen before. Follow it up, explore all around it, and before you know it, you will have something worth thinking about to occupy your mind. All really big discoveries are the result of thought."

Alexander Graham Bell

Contents

1 Introduction **1**

1 The Randall-Sundrum model 6

2 Flat brane multigravity models in five dimensions **8**

1 The $'' + -+''$ bigravity model 8

2 The GRS model and its relation to the $'' + -+''$ bigravity model 18

2.1 Spectrum Interpolation 22

2.2 Multigravity 24

3 The Crystal Universe Model 28

4 Propagator structure 34

3 Radions and dilaton **37**

1 The general three three-brane system 38

2 Effective action 39

3 Scalars Kinetic Energy 42

3.1 The compact case 42

3.2 The non-compact limit 43

4 Gravity on the branes 45

5 Review of sum rules 51

6 Dilaton in warped backgrounds 53

4 AdS_4 brane multigravity models in five dimensions **56**

1 The $'' + +''$ bigravity model 57

2 Graviton propagator in dS_4 and AdS_4 space 67
3 Discussion 72

5 Multigravity in six dimensions **75**
1 Single brane models in six dimensions 76
1.1 The minimal single brane model 76
1.2 The generalized single brane model 78
2 Bigravity in six dimensions 81
2.1 The conifold model 82
2.2 The non-singular model 86
3 Multigravity in six dimensions 88
3.1 Quasi-localized gravity 88
3.2 The crystal universe model 91
4 The weaker energy condition 92

6 Conclusions **94**

A Effective action for graviton-radion-dilaton system **96**

B Ricci tensor and scalar for the dilaton perturbation **99**

C Equations of motion in six dimensions **101**

Bibliography **103**

Publications

The second chapter is based on works published in Nucl. Phys. B **584**, 313 (2000) and in Nucl. Phys. B **595**, 225 (2001). The third chapter has been published in Phys. Lett. B **505**, 231 (2001) and in Nucl. Phys. B **625**, 179 (2002). The fourth chapter is based on works published in Phys. Lett. B **501**, 140 (2001) and in Phys. Lett. B **503**, 173 (2001). Finally, the contents of the fifth chapter are published in Phys. Rev. D **64**, 124014 (2001).

Acknowledgements

First and foremost, I would like to express my deepest gratitude to the people that have guided me in the paths of Theoretical Physics the past three years. I am indebted to my supervisor, collaborator and friend Ian I. Kogan for our countless stimulating discussions, his continuous encouragement and his limitless enthusiasm about physics that inspired me during my stay in Oxford. I am equally grateful to Graham G. Ross for giving me the privilege to collaborate with him the past two years and for generously sharing with me time and ideas that have stamped my short academic life. Without any doubt I was more than fortunate to have such teachers.

Secondly, I would like to thank all the people with whom I have had rewarding collaborations and discussions. In particular, I thank Thibault Damour for very illuminating discussions and for crucial suggestions on the manuscript of this thesis. I would like to thank Stavros Mouslopoulos for collaborating the past years and for our numerous joint publications. I am also grateful to Luigi Pilo and José Santiago for our common work and for very fruitful discussions. Moreover, I would like to thank Panagiota Kanti for valuable correspondence during the past six months. Furthermore, I want to thank all the people from Theoretical Physics for our endless conversations about physics and not only, especially my colleagues in my offices 1.12 and 6.1.

On the practical side of the things, I would like to thank the Hellenic State Scholarship Foundation (IKY) for supporting my studies. I am also grateful to the Department of Theoretical Physics and to Keble College for the travel grands that made it possible for me to participate to various schools, conferences and workshops the last three years. I also would like to thank the coordinators of the "Beyond the Standard Model" and "String Phenomenology" UK networks for giving me the opportunity to attend and talk in the meetings they organized.

During these three years in Oxford I have met a lot of good friends that have made my stay particularly enjoyable. Thanks a lot Alejandro, Alex, Austin, Bayram, Dan, Eleni, Emiliano, Eric, Eva, Fabien, Geza, Katerina, Liliana, Marianna, Mário, Mark, Martin, Martin, Michael, Nuno, Peter, Peter, Ramon, Riccardo, Robert and Shinsuke.

At last but not least, I am grateful to my parents Eleni and Yiannis, my sister Maria and my brother-in-law Tassos for their continuous encouragement, support and above all their love.

Chapter 1

Introduction

There has been a lot of attention over the past few years in theories where the Standard Model (SM) fields are confined on a three dimensional dynamical hypersurface, a three-brane, in a higher than four dimensional spacetime. This scenario is widely known as the brane-world conjecture. This idea dates back to the early eighties when independently Akama [1] and Rubakov and Shaposhnikov [2] suggested models of our Universe as a topological defect and later Visser [3] and Squires [4] described how particles can be gravitationally trapped on the brane. This conjecture has been subject of renewed interest in the recent years with the realization that such structures are common in string theory where the presence of D-branes [5] where open stings end is ubiquitous.

We can generally divide the recent brane-world models into two classes. The first one pioneered by Antoniadis, Arkani-Hamed, Dimopoulos and Dvali [6, 7, 8] assumes that we live on a 3-brane in a $(3+1+n)$-dimensional spacetime with factorizable geometry along the extra dimensions. The higher dimensional Planck scale M is then related to the 4D Planck scale by $M_{\rm Pl}^2 = M^{n+2} V_n$ where V_n is the compactification volume. Taking the size of the new n dimensions to be sufficiently large and identifying the $(4+n)$-Planck scale with the electroweak scale (TeV), a hierarchy between the electroweak and the Planck scale is generated. This gives an alternative to technicolour or low energy supersymmetry resolution to the hierarchy problem. Experimental and astrophysical constraints demand that $n \geq 2$, $M \gtrsim 30$TeV and allow new dimensions even of submillimeter size. However, a new hierarchy must now be explained, namely the ratio of the large compactification radius to the electroweak scale. Even for six extra dimensions this must be greater than 10^5.

The second class of models which attracted equally considerable attention were the ones involving non-factorizable geometry along the extra dimensions and have quite distinct phenomenology from the factorizable geometry case. These models come under the name of warped brane constructions and are motivated by heterotic M-theory [9,10,11] and its five dimensional reduction [12,13]. In the context of these models, one can localize gravity on the brane world having four dimensional gravity even with one extra dimension of infinite extent [14,15] (RS2), or can generate an exponential scale hierarchy with a compact two brane model of Planck scale volume as it was done in the Randall-Sundrum (RS1) model [16], providing a novel geometrical resolution of the hierarchy problem. The scale hierarchy is generated by an exponential with respect to the extra dimension warp factor multiplying the Minkowski metric. We will briefly review in more details this model at the end of this introduction since we will concentrate on warped brane-world models.

In this thesis we will discuss another interesting possibility which arises from warped brane-world constructions, the scenario of multigravity. The proposal of multigravity is that the gravitational interaction that we feel in our Universe is the result of the combination of massless and massive gravitons or even of massive gravitons only. The extra dimensional models give us an infinite tower of massive Kaluza-Klein (KK) gravitons after the process of dimensional reduction. However, one should keep in mind that Newton's inverse square gravitational law is very well tested for distances that range from the submillimeter regime up to at most 10^{26}cm that corresponds to 1% of the size of the observable Universe (see Fig.1.1). Therefore, the massive KK gravitons that will give rise to multigravity should have Compton wavelengths bigger than 10^{26}cm. Additionally, the remaining of the KK tower should either have wavelengths entirely below the submillimeter scale or if they span the whole region they should have very suppressed couplings. It is more than obvious that these requirements are non trivial and cannot be realized in simple KK models since the equidistant spectrum that arises in the typical case will be in contradiction with observations. The aim of this thesis is to demonstrate how one could take such patterns in warped brane-world models. The interesting feature of these multigravity models is that gravity is modified at large scales where one starts to be sensitive to the graviton's mass. We should note at this point that this is the first time that modification of gravity at cosmological scales is triggered by microphysics considerations which have to do with the structure of the extra dimensional space of usually Planck scale volume. These scenarios are in principle testable by the Cosmic Microwave Background (CMB) power spectrum [17,18], gravitational lensing [19] and gravitational wave [20] observations.

We will firstly present multigravity models which involve only flat branes in five dimensions. We will begin our presentation by the $''+-+''$ bigravity model [21] where, apart from the graviton zero mode, an anomalously light graviton KK state is present. This gives the possibility that gravity at intermediate scales ($1\text{mm} < r < 10^{26}\text{cm}$) is mediated by *both* the massless graviton *and* the ultralight state, leading to the terminology "bigravity". At large scales were the Yukawa suppression of the massive state



Figure 1.1: Our current limit of knowledge of gravity.

turns on, the gravitational constant decreases to a value which for example in the symmetric configuration is half of its value at intermediate scales. Similar constructions of freely moving negative tension branes can also give qualitatively different modifications of gravity at large scales. We will briefly review the quasi-localized gravity model of Gregory, Rubakov and Sibiryakov (GRS) [22] which has no normalizable zero mode but gravity at intermediate distances is generated by a "resonance"-like [23, 24] coupling of the lowest part of the continuum of the KK spectrum. In this construction, the gravitational law at large scales changes from four dimensional to five dimensional. Although the bigravity and the quasi-localized gravity models look quite different, they are connected through a more general $''+--+''$ model [25, 26]. We will explicitly show how the spectrum and the gravitational law interpolate between the ones obtained in the above cases. The change of the nature of the gravitational law is observed also in the the crystalline universe model [26] which we will discuss afterwards. In this case gravity at intermediate distances is generated by the first low lying band of KK states and at large scales gives again five dimensional law of gravity. For an alternative way of modifying gravity at ultralarge distances see [27, 28]. At this point we will present a potential danger arising from the fact that massive gravitons have extra polarization states in comparison with the massless one. This is clearly seen by the difference in the tensor structure of the massive and massless spin-2 propagators in flat space (van Dan-Veltman-Zakharov discontinuity) [29, 30]. Observationally this implies that the massive graviton proposal disagrees with well tested predictions of Einsteinian gravity [24].

So far our discussion has considered the gravitational force due to tensorial gravitational modes. However, in the gravity sector there exist moduli corresponding to the fluctuation(s) of the brane(s) or the size of the extra dimension, the radion(s) [32, 33] and the dilaton[1] [34, 35] respectively, which change the phenomenology of these models. This will be the subject of chapter 3 where we will derive the effective action describing these excitations [33]. For this purpose we will firstly consider a general three three-brane model with arbitrary bulk cosmological constants between the branes. We will calculate the kinetic energy for the moduli of the system and see that the freely moving (*i.e.* not on an orbifold fixed point) negative tension branes have wrong sign kinetic term [31, 32, 33]. This plays an important role in the context of multigravity models where this ghost radion(s) cancel the troublesome extra polarization states of the massive gravitons restoring the phenomenologically favoured tensorial structure of the graviton propagator [32, 37, 36]. At large distances the ghost radion(s) dominate giving scalar antigravity [37] in all the above constructions instead of a reduction of the Newton's constant or a change of the nature of the gravitational law. However, the very fact that there are physical ghost(s) in the spectrum makes these models unacceptable. Note that the essential characteristic of all these models is the bounce of the warp factor generated by the presence of the moving negative tension branes. This association between the bounce in the warp factor and presence of negative tension branes is an unavoidable feature in flat (five dimensional) brane models and is possibly linked to the fact that the weaker energy condition is violated at the position of the bounce [38, 39]. In the remaining of the chapter we will discuss the dilaton for backgrounds with identical four-geometries and with bulk dynamics involving arbitrary scalar fields minimally coupled to gravity [40]. In order to do this we will briefly review the brane-world sum rules derived by Gibbons, Kallosh and Linde [41]. Then we will find a generic formula relating the dilaton mass with the sum of the tensions of the branes and the curvature of the four-geometries, from which we can see immediately the necessary condition for stabilizing the overall size of the system.

---

[1] This is not to be confused with the dilaton field that is present in the spectrum of closed strings.

The flat multigravity models as we have already said suffer from the presence of ghost scalar mode(s) which renders the theory unacceptable. In chapter 4 we will present a way out of this difficulty by considering AdS_4 branes instead of flat branes. In this case the warp factor has a $''$cosh$''$ form which naturally generates a bounce without any need of floating negative tension branes. Indeed, it is straightforward to replicate the bigravity and the crystalline model in this framework. The bigravity model is converted to a $''++''$ model [42] and the crystal model to an infinite array of $''+''$ branes. By sending the second brane in the $''++''$ model to infinity, one obtains the locally localized model of [43, 44, 45] where gravity is mediated only by the ultralight state since the zero mode becomes non-normalizable. It is interesting to note how these models confront the relativistic predictions of Einsteinian gravity since there are no ghost radion(s) to restore the graviton propagator structure to the phenomenologically favoured one. We will show that in $(A)dS$ space the van Dam-Veltman-Zakharov no-go theorem can be evaded at tree level for the case that $m/H \to 0$, where m is the mass of the graviton and H the "Hubble" of the $(A)dS$ space, as it was shown in [46, 47, 48, 49]. The extra polarizations of the massive gravitons in this case are very weakly coupled and decouple in the massless flat limit. A precise realization of this case is the $'' + +''$ bigravity model. The *AdS* brane models, even though they solve the theoretical difficulty of the ghost radion(s) of the flat brane models, face phenomenological difficulties. The presence of a remnant negative cosmological constant is at odds with observations, and furthermore it turns out that all large scale modifications of gravity predicted by these models, are hidden behind the *AdS* horizon. Furthermore, as we will discuss at the end of the chapter an old result by Vainshtein [50] suggests that the discontinuity of the graviton propagator in flat space is not directly observable by the bending of the light by the Sun as it was claimed in [29, 30] because non-perturbative effects become important and make the massless limit continuous.

In the light of this encouraging result it seems appropriate to look for multigravity models of flat branes without ghost fields. As we have discussed, in five dimensions it is impossible to have flat brane multigravity models without negative tension branes. For this reason, in chapter 5 we will consider models in six dimensions [51]. We explicitly show that the difficulties of the five dimensional models can be evaded in six dimensions. It is possible to construct multigravity models with only flat positive tension branes. The branes which localize gravity in this setup are four-branes, but one of their dimensions is compact, unwarped and of Planck length. Thus, in the low energy limit the spacetime on the brane appears four dimensional. In order that these constructions are realized it is crucial that the tensions and/or the bulk cosmological constant are anisotropic. The five dimensional constructions in this setup come into two types. One of them involves conical singularities at finite proper distance from the four branes. These conical singularities support three-branes which can be of positive tension if one has an angle deficit, of zero tension if one has no angle deficit and of negative tension if one has an angle excess. The other type of the multigravity models has no conical singularity at all. It is crucial to note that there is no five dimensional effective theory for these constructions, otherwise one would get all the problems faced in the five dimensional constructions. The low energy effective theory is directly four dimensional. These six dimensional models give for the first time a theoretically and phenomenologically viable realization of multigravity.

In the final chapter, we summarize the multigravity models that we have presented in this thesis and discuss questions for future investigation.

Throughout this thesis we use the mostly plus metric convention with signature being $(-,+,+,+,\dots)$ and we use the following convention for the Riemann tensor $R^K_{\Lambda MN} = \partial_M \Gamma^K_{\Lambda N} - \partial_N \Gamma^K_{\Lambda M} + \Gamma^H_{\Lambda N}\Gamma^K_{MH} - \Gamma^H_{\Lambda M}\Gamma^K_{NH}$ where the Christoffel symbols are $\Gamma^K_{MN} = \frac{1}{2} g^{K\Lambda} \left(\partial_M g_{\Lambda N} + \partial_N g_{\Lambda M} - \partial_\Lambda g_{MN}\right)$. We work in the units where $\hbar = c = 1$. The four dimensional Planck mass and Newton's constant are related as $G_N^{-1} = 32\pi M_{\rm Pl}^2$.



Figure 1.2: The Randall-Sundrum model with one ″+″ and one ″−″ brane at the fixed points of a S^1/Z_2 orbifold.

1 The Randall-Sundrum model

Since, throughout this thesis we will study warped models it is instructive to briefly present the Randall-Sundrum model. This construction consists of two parallel three-branes sitting on the fixed points of an S^1/Z_2 orbifold in five dimensional spacetime which is essentially a slice of AdS_5 (see Fig.1.2). The negative bulk cosmological constant and the tensions of the branes are fine tuned in such a way that the four-dimensional slices of the five-dimensional spacetime, and thus the two branes, are flat. The metric that describes this configuration has the characteristic warped form:

$$ds^2 = e^{-2k|y|}\eta_{\mu\nu}dx^\mu dx^\nu + dy^2 \tag{1.1}$$

If one now considers a scalar field, in particular the SM Higgs, confined on one of the two branes, then the minimal coupling to the metric will result to an action:

$$\mathcal{S}_H = \int d^4x \underbrace{\sqrt{-g_{ind}}}_{e^{-4k|y_i|}} \left\{ \underbrace{-\ g^{\mu\nu}_{ind}}_{-e^{2k|y_i|}\eta^{\mu\nu}} D_\mu H^\dagger D_\nu H - \lambda \left(H^2 - v_0^2\right)^2 \right\} \tag{1.2}$$

where $g^{ind}_{\mu\nu}$ is the induced metric on the brane at position y_i. After a field rescaling $H \to e^{k|y_i|}H$ which brings the kinetic term to be canonical, the action reads:

$$\mathcal{S}_H = \int d^4x \left\{ -\eta^{\mu\nu} D_\mu H^\dagger D_\nu H - \lambda \left(H^2 - v_0^2 e^{-2k|y_i|}\right)^2 \right\} \tag{1.3}$$

which clearly shows that the vacuum expectation value (vev) of the Higgs is given as:

$$v = v_0 e^{-k|y_i|} \tag{1.4}$$



Figure 1.3: The graviton (blue line), first (red line) and second (red dashed line) KK states wavefunctions in the RS1 model. On the right the pattern of the spectrum is sketched.

Assuming that all five dimensional parameters and thus v_0 are of the order of the Planck scale, then a Higgs on the first brane will have a vev of the order of the Planck scale, whereas a Higgs on the second brane will have exponentially reduced vev. It was exactly this generation of exponentially different mass scales that was utilized in the Randall-Sundrum scenario to resolve the Planck hierarchy problem by placing the SM on the second brane (TeV brane). A compactification radius of only some 35 times larger than the Planck length was sufficient to bring the Higgs vev to the TeV scale.

The phenomenology of this model has been extensively explored over the past years (see [52] for a comprehensive account). In the TeV hierarchy scenario, the KK tower of the spin-2 graviton resonances starts from the TeV scale with approximately TeV spacing between them giving rise to characteristic signals in high energy colliders. The graviton and KK states wavefunctions (*i.e.* their profiles along the extra dimension) are shown in Fig.1.3. The massless zero mode wavefunction follows the warp factor and is localized on the positive tension brane, while the wavefunctions of the remaining tower are suppressed on the positive tension brane and enhanced on the negative tension one. We will return to this picture later and compare it with the considerably different pattern that we get in the multigravity models.

Chapter 2

Flat brane multigravity models in five dimensions

In this chapter we will see how we can realize the multigravity scenario, as it has been described in the introduction, with flat multi-brane configurations in five dimensions. We will see that these models are infested with the presence of freely moving negative tension branes. We will discuss the problems arising from the negative tension branes in the following chapter. Here, we will concentrate on the gravitational potential coming from the tensor fluctuations of the background metric and discuss the general characteristics of the multigravity models. Although flat brane multigravity scenarios have difficulties, they are very easy to handle toy models and can be used to understand physics of more complicated but realistic multigravity models.

In the following we will discuss the $'' + -+''$ bigravity modes, the GRS model of quasi-localized gravity and the relation between them as well as the crystal multigravity model.

1 The $'' + -+''$ bigravity model

We begin our presentation of flat multigravity models in five dimensions with the $'' + -+''$ bigravity model [21]. This model is a straightforward extension of the RS1 model with one more positive tension brane. In more detail, we have three parallel three-branes in an AdS_5 space with cosmological constant $\Lambda < 0$. The fifth dimension has the geometry of an S^1/Z_2 orbifold and the branes are located at $L_0 = 0$, L_1 and L_2 where L_0 and L_2 are the orbifold fixed points (see Fig.2.1). This model has three parameters, namely L_1, L_2 and Λ. For our present purposes we will concentrate in the symmetric configuration where the intermediate brane lies in the middle of the orbifold, *i.e.* $L_2 = 2L_1 \equiv 2l$, leaving two parameters l, Λ. Firstly, we consider the branes having no matter on them in order to find a suitable vacuum solution. The



Figure 2.1: The ″ + −+″ bigravity model with two ″+″ branes at the fixed points of a S^1/Z_2 orbifold and a ″−″ brane in between.

action of this setup is:

$$\mathcal{S}_G = \int d^4x \int_{-2l}^{2l} dy \sqrt{-G}\{2M^3 R - \Lambda\} - \sum_i \int_{y=L_i} d^4x V_i \sqrt{-\hat{G}^{(i)}} \tag{2.1}$$

where $\hat{G}^{(i)}_{\mu\nu}$ is the induced metric on the branes, V_i their tensions and M the five dimensional fundamental scale. The integration is performed throughout the orbifold, so there is no need to introduce Gibbons-Hawking boundary terms. The Einstein equations that arise from this action are:

$$R_{MN} - \frac{1}{2} G_{MN} R = -\frac{1}{4M^3}\left(\Lambda G_{MN} + \sum_i V_i \frac{\sqrt{-\hat{G}^{(i)}}}{\sqrt{-G}} \hat{G}^{(i)}_{\mu\nu} \delta^\mu_M \delta^\nu_N \delta(y - L_i)\right) \tag{2.2}$$

At this point we demand that our metric respects four dimensional Poincaré invariance. The metric ansatz with this property is the following:

$$ds^2 = e^{-2\sigma(y)} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2 \tag{2.3}$$

Here the warp function $\sigma(y)$ is essentially a conformal factor that rescales the four dimensional component of the metric. A straightforward calculation gives us the following differential equations for $\sigma(y)$:

$$(\sigma')^2 = k^2 \tag{2.4}$$

$$\sigma'' = \sum_i \frac{V_i}{12M^3} \delta(y - L_i) \tag{2.5}$$

where $k=\sqrt{\frac{-\Lambda}{24M^3}}$ is a measure of the curvature of the bulk.

The solution of these equations consistent with the orbifold geometry is precisely:

$$\sigma(y)=k\left\{l-||y|-l|\right\} \tag{2.6}$$

with the additional requirement that $V_0=V_2=-V_1\equiv V$. Furthermore, four dimensional Poincaré invariance like in the RS1 model requires the fine tuning relation:

$$V=-\frac{\Lambda}{k} \tag{2.7}$$

If we consider massless fluctuations of this vacuum metric as in [16] and then integrate over the fifth dimension, we find the four dimensional fundamental scale[1] M_*, the five dimensional fundamental scale M and the curvature of the bulk k are related as:

$$M_*^2=\frac{2M^3}{k}\left[1-e^{-2kl}\right] \tag{2.8}$$

We do not identify the mass scale M_* with the four dimensional Planck scale for reasons which will be obvious in the following. The above formula tells us that for large enough kl the three mass scales M_*, M, k can be taken to be of the same order. Thus we take $k\sim\mathcal{O}(M)$ in order not to introduce a hierarchy, with the additional restriction $k<M$ so that the bulk curvature is small compared to the five dimensional Planck scale and we can trust our solution. Furthermore, if we put matter on the branes all the physical masses m on the brane will be related to the fundamental mass parameters $m_0\sim M$ of the five dimensional theory by the conformal warp factor:

$$m=e^{-\sigma(L_i)}m_0 \tag{2.9}$$

Thus, in the symmetric configuration which we are examining, the mass scales on the first and third brane will be of the order of the fundamental scale M while on the intermediate negative tension brane they will be exponentially suppressed. It is obvious that in an asymmetric configuration we could obtain an electroweak scale hierarchy a la Randall-Sundrum even on a positive tension brane. For this case we would only need $2L_1-L_2\approx 35k^{-1}$ in order to get $e^{-\sigma(L_3)}\sim\mathcal{O}\left(10^{-15}\right)$ (see [21,53]).

To determine the phenomenology of the model we need to know the KK spectrum that follows from the dimensional reduction. This is determined by considering the linear fluctuations of the metric of the form:

$$ds^2=\left[e^{-2\sigma(y)}\eta_{\mu\nu}+h_{\mu\nu}(x,y)\right]dx^\mu dx^\nu+dy^2 \tag{2.10}$$

---

[1] The factor M_* multiplies the four dimensional Ricci scalar in the zero mode truncated Lagrangian.

Here we have ignored the dilaton and radion modes which will be the subject of the following chapter. We expand the field $h_{\mu\nu}(x,y)$ in graviton and KK states plane waves according to:

$$h_{\mu\nu}(x,y)=\sum_{n=0}^{\infty}h_{\mu\nu}^{(n)}(x)\Psi^{(n)}(y) \tag{2.11}$$

where $\left(\partial_\kappa\partial^\kappa-m_n^2\right)h_{\mu\nu}^{(n)}=0$ with $\partial^\alpha h_{\alpha\beta}^{(n)}=h^{(n)\alpha}_{\ \ \ \alpha}=0$. The latter conditions for the zero mode correspond to a choice of gauge fixing. The function $\Psi^{(n)}(y)$ will obey the following second order differential equation:

$$\left\{-\frac{1}{2}\partial_y^2+2k^2-2k\left[\delta(y)+\delta(y-2l)-\delta(y-l)-\delta(y+l)\right]\right\}\Psi^{(n)}(y)=\frac{m_n^2}{2}e^{2\sigma}\Psi^{(n)}(y) \tag{2.12}$$

After a change of variables and a redefinition of the wavefunction the above equation reduces to an ordinary Schrödinger-type equation:

$$\left\{-\frac{1}{2}\partial_z^2+V(z)\right\}\hat{\Psi}^{(n)}(z)=\frac{m_n^2}{2}\hat{\Psi}^{(n)}(z) \tag{2.13}$$

with potential:

$$V(z)=\frac{15k^2}{8[g(z)]^2}-\frac{3k}{2g(z)}\left[\delta(z)+\delta(z-2z_l)-\delta(z-z_l)-\delta(z+z_l)\right] \tag{2.14}$$

The new variables and wavefunction in the above equation are defined as:

$$z\equiv\mathrm{sgn}(y)\begin{cases}\frac{2e^{kl}-e^{2kl-k|y|}-1}{k} & ,l\leq|y|\leq 2l\\ \frac{e^{k|y|}-1}{k} & ,|y|\leq l\end{cases} \tag{2.15}$$

$$\hat{\Psi}^{(n)}(z)\equiv\Psi^{(n)}(y)e^{\sigma/2} \tag{2.16}$$

and the function $g(z)$ as $g(z)\equiv k\left\{z_l-||z|-z_l|\right\}+1$, where $z_l=z(l)$. The change of variables has been chosen so that there are no first derivative terms in the differential equation. Furthermore, in this coordinate system it can be easily seen that the vacuum metric takes the conformally flat form:

$$ds^2=\frac{1}{g(z)^2}\left(\eta_{\mu\nu}dx^\mu dx^\nu+dz^2\right) \tag{2.17}$$

This is a quantum mechanical problem with δ-function potentials and an extra $1/g^2$



Figure 2.2: The potential $V(z)$ in the $''+-+''$ bigravity model. The δ-function wells correspond to $''+''$ branes, while the δ-function barriers to $''-''$ branes.

smoothing term (due to the AdS geometry) that gives the potential a double "volcano" form (see Fig.2.2). An interesting characteristic of this Schrödinger equation is that it has a "supersymmetric" form which guarantees that it has no tachyonic modes:

$$AA^{\dagger}\hat{\Psi}^{(n)}(z) = m_n^2\hat{\Psi}^{(n)}(z) \quad \text{with} \quad A = \frac{d}{dz} - \frac{3}{2}\frac{\partial_z g}{g} \tag{2.18}$$

Since the system is compact, this differential equation always gives rise to a massless zero mode which reflects the fact that Poincaré invariance is preserved in the four dimensional spacetime. Its wavefunction is given by:

$$\hat{\Psi}^{(0)} = \frac{A}{[g(z)]^{3/2}} \tag{2.19}$$

The normalization factor A is determined by the requirement $\int_{-2z_l}^{2z_l} dz \left[\hat{\Psi}^{(0)}(z)\right]^2 = 1$, chosen so that we get the standard form of the Fierz-Pauli Lagrangian.

In the symmetric case that we are examining, the potential, and thus the zero mode wavefunction, is symmetric with respect to the second brane. This immediately implies that the next level will have a wavefunction antisymmetric with respect to the second brane. This is exactly what happens. The solution for the graviton KK modes is given in terms of Bessel functions. For y lying in the regions $\mathbf{A} \equiv [0, l]$ and $\mathbf{B} \equiv [l, 2l]$, we have:

$$\hat{\Psi}^{(n)} \left\{ \begin{array}{c} \mathbf{A} \\ \mathbf{B} \end{array} \right\} = \sqrt{\frac{g(z)}{k}} \left[\left\{ \begin{array}{c} A_1 \\ B_1 \end{array} \right\} J_2\left(\frac{m_n}{k} g(z)\right) + \left\{ \begin{array}{c} A_2 \\ B_2 \end{array} \right\} Y_2\left(\frac{m_n}{k} g(z)\right) \right] \tag{2.20}$$

The boundary conditions (one for the continuity of the wavefunction at z_1 and three for the discontinuity of its first derivative at 0, z_1, z_2) result in a 4×4 homogeneous linear system which, in order to have a non-trivial solution, should have vanishing determinant:

$$\left| \begin{array}{cccc} J_1\left(\frac{m}{k}\right) & Y_1\left(\frac{m}{k}\right) & 0 & 0 \\ 0 & 0 & J_1\left(\frac{m}{k}\right) & Y_1\left(\frac{m}{k}\right) \\ J_1\left(\frac{m}{k} g(z_l)\right) & Y_1\left(\frac{m}{k} g(z_l)\right) & J_1\left(\frac{m}{k} g(z_l)\right) & Y_1\left(\frac{m}{k} g(z_l)\right) \\ J_2\left(\frac{m}{k} g(z_l)\right) & Y_2\left(\frac{m}{k} g(z_l)\right) & -J_2\left(\frac{m}{k} g(z_l)\right) & -Y_2\left(\frac{m}{k} g(z_l)\right) \end{array} \right| = 0 \tag{2.21}$$

Here we have suppressed the subscript n on the masses m_n. This is essentially the mass quantization condition which gives the spectrum of the KK states. For each mass we can then determine the wavefunction with the same normalization as the zero mode.

The exact calculation of the wavefunction of the first graviton KK state verifies our previously stated intuition. Another way to understand the form of the wavefunction is that since we have two delta functions in the potential each of which supports a bound state, there should be a massive KK state that resembles the zero mode. Indeed, if one calculates the wavefunctions of the higher levels it is evident that the first KK state has a special characteristic. It is the only one that is peaked in the positive tension branes, whereas all the higher levels are suppressed on the positive tension branes and are spread in the bulk (see Fig.2.3). This is in contrast to the RS1 case where the wavefunctions of all the KK states are spread in the bulk (see Fig.1.3). Moreover, since the difference of the modulus of the wavefunctions of the first special KK state and the zero mode comes from a very small region of the y-space around the position of the minimum of the warp factor, we expect its mass to be anomalously light in comparison to the masses of the remaining of the KK tower.

This is a generic characteristic of potentials which have the shape of a bounce. In five dimensions with flat branes the only possibility of obtaining a bounce is by introducing negative tension branes. This is because the warp factors generated by flat positive tension branes always have the form of a falling exponential. The only way



Figure 2.3: The graviton (blue line), first (red line) and second (green line) KK states wavefunctions in the symmetric $''+-+''$ model. The wavefunctions of the zero mode and the second KK state have discontinuities on the $''-''$ branes. On the right the pattern of the spectrum is sketched.

of pasting two warp factors which are falling exponentials is through a discontinuity generated by a negative tension brane.

If we define $x = kl$, as the distance between the negative and positive branes measured in units of k, then for $x \gg 1$ we may obtain a reliable approximation to its mass by using the first terms of the Bessel power series. Then we find that the first KK mode has mass given by:

$$m_1 = 2\sqrt{2}ke^{-2x} \tag{2.22}$$

The masses of the other KK states are found to depend in a different way on the parameter x. Analytically we find that for $x \gg 1$ the masses are:

$$m_{n+1} = \xi_n ke^{-x} \quad , \quad n = 1, 2, \dots \tag{2.23}$$

where ξ_{2i+1} is the $(i+1)$-th root of $J_1(x)$ $(i = 0, 1, 2, \dots)$ and ξ_{2i} is the i-th root of $J_2(x)$ $(i = 1, 2, \dots)$. Using the asymptotic values of the Bessel roots, we find that the level splitting for high n levels approaches a constant value:

$$\Delta m = \frac{\pi}{2}ke^{-x} \tag{2.24}$$

The coupling also of the first KK state to matter on the branes is very different from the one of the remaining tower. To see this, we consider a minimally coupled to gravity matter Lagrangian $\mathcal{L}_m(\Phi_j, \hat{G}^{(i)})$, where we denote with Φ_j the matter fields living on the branes. The total action of the system is the sum of the gravity action (2.1) plus the matter action:

$$\mathcal{S} = \mathcal{S}_G + \sum_i \int d^4x dy \sqrt{-\hat{G}^{(i)}} \mathcal{L}_m(\Phi_j, \hat{G}^{(i)}) \tag{2.25}$$

The effective four dimensional Lagrangian taking into account only the graviton perturbations, has the following form:

$$\mathcal{L} = \mathcal{L}_m(\Phi_j^c, \eta) + 2M^3 \sum_n \mathcal{L}_{PF}(h^{(n)}(x)) - \sum_i \sum_n \frac{e^{2\sigma(L_i)}\Psi^{(n)}(L_i)}{2} h_{\mu\nu}^{(n)}(x) T_{(i)}^{\mu\nu} \tag{2.26}$$

where $\mathcal{L}_{PF}$ is the Pauli-Fierz Lagrangian for the n-level graviton mode. The matter fields have been rescaled $\Phi_j \rightarrow \Phi_j^c$, to make them canonically normalized and the energy momentum tensor $T_{(i)}^{\mu\nu}$ for matter localized on the i-th brane is defined with respect to the rescaled fields Φ_j^c. By construction the induced background metric on the branes is the flat four dimensional Minkowski metric ($\eta_{\mu\nu}$). Finally, defining the canonically normalized graviton fields:

$$\bar{h}_{\mu\nu}^{(n)}(x) = \sqrt{2M^3} h_{\mu\nu}^{(n)}(x) \tag{2.27}$$

the Lagrangian reads:

$$\mathcal{L} = \mathcal{L}_m(\Phi_j^c, \eta) + \sum_n \mathcal{L}_{PF}(\bar{h}^{(n)}(x)) - \sum_i \sum_n \frac{\hat{\Psi}^{(n)}(z_i)\, g(z_i)^{3/2}}{2\sqrt{2M^3}} \bar{h}_{\mu\nu}^{(n)}(x)\, T_{(i)}^{\mu\nu} \tag{2.28}$$

Thus, the gravitons have the following couplings to matter on the positive tension branes:

$$a_n = \frac{\hat{\Psi}^{(n)}(z_+)\, g(z_+)^{3/2}}{2\sqrt{2M^3}} \tag{2.29}$$

In more detail we find that the zero mode and the first ultralight KK state couple with the same strength on the positive tension branes:

$$a_0 = a_1 = \frac{1}{2\sqrt{2}M_*} \tag{2.30}$$

while the coupling of the remaining tower is exponentially suppressed:

$$a_{n+1} = \frac{1}{2\sqrt{2}M_*} \frac{e^{-x}}{\sqrt{J_1^2\left(\xi_n\right) + J_2^2\left(\xi_n\right)}} \qquad n = 1, 2, 3, \dots \tag{2.31}$$

The reason for the strong coupling of the first KK state in comparison with the remaining tower is readily understood because, being dominantly a bound state of the volcano potential on the positive tension brane, it is largely localized on it.

Let us now demonstrate how the bigravity scenario can be realized in this model. We have already seen from (2.22) and (2.23) that, for large x, the lightest KK mode splits off from the remaining tower. This allows the possibility that the Newtonian gravity that we experience at intermediate distances ($1\text{mm} < r < 10^{26}\text{cm}$), is due to *both* the massless graviton *and* the ultralight KK state. The gravitational potential is computed by the tree level exchange diagrams of the massless graviton and KK states which in the Newtonian limit is:

$$V(r) = -\sum_{n=0}^{N_\Lambda} \frac{a_n^2}{4\pi} \frac{e^{-m_n r}}{r} \tag{2.32}$$

where a_n is the coupling (2.29) and $n = 0$ accounts for the massless graviton. The summation stops at some very high level N_Λ with mass of the order of the cutoff scale $\sim M$. In the bigravity scenario, at intermediate distances $m_2^{-1} \ll r \ll m_1^{-1}$, the first KK state and the massless graviton contribute equally to the gravitational potential:

$$V_{id}(r) = -\frac{1}{32\pi M_*^2}\left(\frac{1}{r} + \frac{1}{r}\right) = -\frac{G_N}{r} \tag{2.33}$$

where $G_N \equiv \frac{1}{16\pi M_*^2}$ ($M_{\text{Pl}} = \frac{M_*}{\sqrt{2}}$). For long distances $r \gtrsim m_1^{-1}$ the Yukawa suppression effectively reduces gravity to half its strength:

$$V_{ld}(r) = -\frac{1}{32\pi M_*^2}\left(\frac{1}{r} + \frac{0}{r}\right) = -\frac{G_N}{2}\frac{1}{r} \tag{2.34}$$

Astronomical constraints and the requirement of the observability of this effect demand that for $k \sim M_{\text{Pl}}$ we should have x in the region $65-70$. The observational and experimental constrains in the context of the graviton KK modes are $m < 10^{-31}\text{eV}$ or $m > 10^{-4}\text{eV}$ for modes with Planck suppressed coupling. If we choose $m_1 \approx 10^{-31}\text{eV}$, from (2.22) and (2.23) we find $m_2 \approx 10^{-2}\text{eV}$ which comfortably satisfies the Cavendish bound (see Fig.2.4).


0
10μm
0.2mm
Excluded by Observational Data
and by the Cavendish Experiments
10^{26}cm
r
m_2^{-1}
m_1^{-1}

Figure 2.4: Exclusion regions for the bigravity case and correlation of the first two KK states.

However, the Cavendish bound in this case is not even applicable because the coupling of the remaining KK tower is exponentially smaller than the usual Planck suppressed coupling. To see this let us calculate the correction to the gravitational law at distances $r \lesssim m_2^{-1}$. The Yukawa interactions of the remaining KK states then turn on and will give rise to a short distance correction. This can be evaluated by using the asymptotic expression of the Bessel functions in (2.31) since we are dealing with large x and summing over a very dense spectrum, giving:

$$\Delta V_{sd}(r) = -\frac{G_N}{k}\sum_{n=2}^{N_\Lambda}\frac{k\pi}{2e^x}\,\frac{m_n}{2k}\,\frac{e^{-m_n r}}{r} \tag{2.35}$$

At this point we exploit the fact that the spectrum is nearly continuum above m_2 and turn the sum to an integral with the first factor in (2.35) being the integration measure, *i.e.* $\sum \frac{k\pi}{2e^x} = \sum \Delta m \to \int dm$ from (2.24). Moreover, we can extend the integration to infinity because, due to the exponential suppression of the integrand, the integral saturates very quickly and thus the integration over the region of very large masses is irrelevant. The resulting potential is now:

$$\Delta V_{sd}(r) = -\frac{G_N}{k}\int_{m_2}^{\infty} dm\,\frac{m}{2k}\,\frac{e^{-mr}}{r} \tag{2.36}$$

The integration is easily performed and gives:

$$\Delta V_{sd}(r) = -\frac{G_N}{2r}\,\frac{1+m_2 r}{(kr)^2}\,e^{-m_2 r} \tag{2.37}$$

We see these short distance corrections are significant only at Planck scale distances $\sim k^{-1}$. Thus, the lower bound of the Cavendish experiments is not important in the symmetric configuration simply because although the low lying KK states have Compton wavelength at the millimeter scale, they are however very weakly coupled.

Finally, let us comment about the realization of the bigravity scenario in asymmet-



Figure 2.5: The GRS model with one $''+''$ brane at the fixed point of an R^1/Z_2 orbifold and a $''-1/2''$ brane at a distance $x = kL$. The curvature of the bulk between the $''+''$ and $''-1/2''$ branes is k and zero in the outside region.

ric configurations where the positive tension branes are not scale equivalent. In this case a whole rich of bigravity scenarios emerge. Depending on which positive brane we live in, we can have either a dominant massless graviton component or a dominant massive graviton component. In the first case the correction to Newtonian gravitational law at large scales will be negligible. In the second case, where the massive KK state is the dominant source of Newtonian gravity, one would see gravity to essentially switch off at large scales.

2 The GRS model and its relation to the $''+-+''$ bigravity model

An other interesting model which also has the feature of predicting modifications of gravity at ultralarge scales is the one by Gregory, Rubakov and Sibiryakov (GRS) [22]. In this section we will briefly review the physics of the GRS model and demonstrate how it is related to the $''+-+''$ bigravity model. The GRS model modifies the RS2 model by adding a negative tension brane of half the tension of the positive tension one at some distance L in the bulk. The space between the positive and negative tension branes is a slice of AdS_5, while in the outside region it is flat (see Fig.2.5). The background metric for this setup has the four dimensional Poincaré invariant form (2.3) with the warp factor being:

$$\sigma(y) = \left\{ \begin{array}{cc} k|y| & ,|y| < L \\ kL & ,|y| > L \end{array} \right. \tag{2.38}$$

This model does not have a normalizable massless graviton because the volume of the system is infinite, but has a continuum of KK states that starts from zero. Despite the absence of a zero mode, it generates four dimensional gravity at intermediate distances due to the “resonance”-like behaviour of the values of the wavefunctions of the KK states continuum at the position of the positive tension brane. In that respect, gravity in this picture appears to be quasi-localized. The coupling for small masses of the KK states has a Breit-Wigner form:

$$a(m)^2 = \frac{\mathcal{A}}{m^2 + (\Gamma/2)^2} \tag{2.39}$$

where $\Gamma = 4ke^{-3x}$ is the width of the resonance. This feature of the coupling gives for intermediate distances $r \ll \Gamma^{-1}$ normal four dimensional gravitational law:

$$V_{id}(r) = -G_N \frac{1}{r} \tag{2.40}$$

where $G_N = \frac{k}{32\pi M^3}$. For long distances $r \gg \Gamma^{-1}$, however, the gravitational law becomes five dimensional:

$$V_{ld}(r) = -\frac{4G_N}{\pi\Gamma}\frac{1}{r^2} \tag{2.41}$$

If we have $L \approx 45k^{-1}$, then $\Gamma^{-1} \sim 10^{26}$cm, so this scenario cannot be excluded by current observations. The short distance corrections, as in the $''+-+''$ model, appear only for $r \lesssim k^{-1}$ and read:

$$\Delta V_{sd}(r) = -\frac{G_N}{2r}\frac{1}{k^2r^2} \tag{2.42}$$

Although the GRS and the $''+-+''$ bigravity models look quite different, they share the key element of freely moving negative tension branes. It was shown in [26] that these two models are the limiting cases of a more general $''+--+''$ multi-brane model that interpolates between them. This model can be readily derived from the bigravity model by cutting in half the intermediate negative tension brane, *i.e.* instead of having one negative tension brane one can take two negative tension branes of half the tension of the original one, and considering flat space between the two new negative tension branes. If the second positive tension brane is taken to infinity together with one of the negative tension branes, we shall get precisely the GRS picture.

Let us discuss the four-brane $''+--+''$ model in more detail. It consists of a five dimensional spacetime with orbifold topology with four parallel three-branes located at $L_0 = 0$, L_1, L_2 and L_3, where L_0 and L_3 are the orbifold fixed points (see Fig.2.6). The bulk cosmological constant Λ is negative between the positive and negative tension branes and zero between the two negative tension branes. The model has four parameters, namely L_1, L_2 and L_3 and Λ. For our present purposes we consider the symmetric configuration, leaving three parameters, l, l_- and Λ where



Figure 2.6: The $'' + - - +''$ configuration with scale equivalent $''+''$ branes. The distance between the $''+''$ and $'' - 1/2''$ branes is $l = L_1 = L_3 - L_2$ while the distance between the $'' - 1/2''$ branes is $l_- = L_2 - L_1$. The curvature of the bulk between the $''+''$ and $'' - 1/2''$ branes is k and zero between the $'' - 1/2''$ branes.

$l \equiv L_1 = L_3 - L_2$ and $l_- \equiv L_2 - L_1$. In the absence of matter the model is described by (2.1) with

$$\Lambda(y) = \begin{cases} 0 & , y \in [L_1, L_2] \\ \Lambda & , y \in [0, L_1] \bigcup [L_2, L_3] \end{cases} \tag{2.43}$$

By considering the ansatz (2.3) the warp function $\sigma(y)$ must satisfy:

$$\sigma'' = \sum_i \frac{V_i}{12M^3}\delta(y - L_i) \tag{2.44}$$

$$(\sigma')^2 = \begin{cases} 0 & , y \in [L_1, L_2] \\ k^2 & , y \in [0, L_1] \bigcup [L_2, L_3] \end{cases} \tag{2.45}$$

where $k = \sqrt{\frac{-\Lambda}{24M^3}}$ is a measure of the bulk curvature. The solution for $y > 0$ is:

$$\sigma(y) = \begin{cases} ky & , y \in [0, L_1] \\ kl & , y \in [L_1, L_2] \\ kl + k(l + l_- - y) & , y \in [L_2, L_3] \end{cases} \tag{2.46}$$

with the additional requirement that $V_0 = V_3 = -2V_1 = -2V_2 \equiv V$. Furthermore, four dimensional Poincaré invariance requires the fine tuning relation:

$$V = -\frac{\Lambda}{k} \tag{2.47}$$

In order to determine the mass spectrum and the couplings of the KK modes we consider linear fluctuations around the background metric as in (2.10). Following the same procedure we find that the function $\hat{\Psi}^{(n)}(z)$ obeys a Schrödinger-like equation (2.13) with potential $V(z)$ of the form:

$$\begin{aligned} V(z) &= \frac{15k^2}{8[g(z)]^2}\left[\theta(z)-\theta(z-z_1)+\theta(z-z_2)-\theta(z-z_3)\right] \\ &\quad -\frac{3k}{2g(z)}\left[\delta(z)-\frac{1}{2}\delta(z-z_1)-\frac{1}{2}\delta(z-z_2)+\delta(z-z_3)\right] \end{aligned} \tag{2.48}$$

The conformal coordinates now are given by:

$$z \equiv \begin{cases} \frac{e^{ky}-1}{k} & , y \in [0, L_1] \\ (y-l)e^{kl}+\frac{e^{kl}-1}{k} & , y \in [L_1, L_2] \\ -\frac{1}{k}e^{2kl+kl_-}e^{-ky}+l_-e^{kl}+\frac{2}{k}e^{kl}-\frac{1}{k} & , y \in [L_2, L_3] \end{cases} \tag{2.49}$$

with

$$g(z) = \begin{cases} kz+1 & , z \in [0, z_1] \\ kz_1+1 & , z \in [z_1, z_2] \\ k(z_2-z)+kz_1+1 & , z \in [z_2, z_3] \end{cases} \tag{2.50}$$

where $z_1 = z(L_1)$, $z_2 = z(L_2)$ and $z_3 = z(L_3)$.

The potential (2.48) again gives rise to a massless graviton zero mode whose wavefunction is given by (2.19) with the same normalization convention. Note, however, that in the limit $l_- \to \infty$ this mode becomes non-normalizable (GRS case). The solution of the Schrödinger equation for the massive KK modes is:

$$\hat{\Psi}^{(n)}\left\{\begin{array}{c} \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \end{array}\right\} = N_n \left\{\begin{array}{c} \sqrt{\frac{g(z)}{k}}\left[\quad Y_2\left(\frac{m_n}{k}g(z)\right)+A\ J_2\left(\frac{m_n}{k}g(z)\right)\right] \\ B_1\cos(m_n z)+B_2\sin(m_n z) \\ \sqrt{\frac{g(z)}{k}}\left[C_1Y_2\left(\frac{m_n}{k}g(z)\right)+C_2J_2\left(\frac{m_n}{k}g(z)\right)\right] \end{array}\right\} \tag{2.51}$$

where $\mathbf{A} = [0, z_1]$, $\mathbf{B} = [z_1, z_2]$, and $\mathbf{C} = [z_2, z_3]$. We observe that the solution in the first and third interval has the same form as in the $''+-+''$ model. The new feature is the second region (flat spacetime). The coefficients that appear in the solution are determined by imposing the boundary conditions and normalizing the wavefunction.

The boundary conditions (two for the continuity of the wavefunction at z_1, z_2 and four for the discontinuity of its first derivative at 0, z_1, z_2 and z_3) result in a 6×6 homogeneous linear system which, in order to have a non-trivial solution, should have vanishing determinant. It is readily reduced to a 4×4 set of equations leading to the quantization condition:

$$\begin{vmatrix} Y_2\left(\frac{m}{k}g\right)-\frac{Y_1\left(\frac{m}{k}\right)}{J_1\left(\frac{m}{k}\right)}J_2\left(\frac{m}{k}g\right) & -\cos(mz_1) & -\sin(mz_1) & 0 \\ Y_1\left(\frac{m}{k}g\right)-\frac{Y_1\left(\frac{m}{k}\right)}{J_1\left(\frac{m}{k}\right)}J_1\left(\frac{m}{k}g\right) & \sin(mz_1) & -\cos(mz_1) & 0 \\ 0 & -\sin(mz_2) & \cos(mz_2) & Y_1\left(\frac{m}{k}g\right)-\frac{Y_1\left(\frac{m}{k}\right)}{J_1\left(\frac{m}{k}\right)}J_1\left(\frac{m}{k}g\right) \\ 0 & \cos(mz_2) & \sin(mz_2) & -Y_2\left(\frac{m}{k}g\right)+\frac{Y_1\left(\frac{m}{k}\right)}{J_1\left(\frac{m}{k}\right)}J_2\left(\frac{m}{k}g\right) \end{vmatrix} = 0 \tag{2.52}$$

with $g=g(z_1)=g(z_2)$. Here we have suppressed the subscript n on the masses m_n.

2.1 Spectrum Interpolation

The above quantization condition provides the mass spectrum of the model. It is convenient to introduce two dimensionless parameters, $x = kl$ and $x_- = kl_-$ (see Fig.2.6) and we work from now on with the set of parameters x, x_- and k. The mass spectrum depends crucially on the distance x_-. We must recover the $''+-+''$ bigravity model spectrum in the limit $x_- \to 0$, and the GRS spectrum in the limit $x_- \to \infty$. From the quantization condition (2.52) it is easy to verify these features and show how the $''+--+''$ spectrum smoothly interpolates between the $''+-+''$ bigravity model and the GRS one. It turns out that the structure of the spectrum has simple x_- and x dependence in three separate regions of the parameter space:

The $''+-+''$ Bigravity Region

For $x_- \lesssim 1$ we find that the mass spectrum is effectively x_--independent given by the approximate form:

$$m_1 = 2\sqrt{2}ke^{-2x} \tag{2.53}$$

$$m_{n+1} = \xi_n k e^{-x} \qquad n = 1,2,3,\ldots \tag{2.54}$$

where ξ_{2i+1} is the $(i+1)$-th root of $J_1(x)$ $(i = 0,1,2,\ldots)$ and ξ_{2i} is the i-th root of $J_2(x)$ $(i = 1,2,3,\ldots)$. As expected the mass spectrum is identical to the one of the $''+-+''$ bigravity model for scale equivalent positive tension branes. The first mass is manifestly singled out from the rest of the KK tower and for large x leads to the possibility of bigravity (see Fig.2.7).



Figure 2.7: The behaviour of the mass of the first five KK states in the three regions of simple x, x_- dependence. The first dot at zero stands for the graviton.

The Saturation Region

For $1 \ll x_- \ll e^{2x}$ we find a simple dependence on x_- given by the approximate analytic form:

$$m_1 = 2k\frac{e^{-2x}}{\sqrt{x_-}} \tag{2.55}$$

$$m_{n+1} = n\pi k\frac{e^{-x}}{x_-} \qquad n = 1, 2, 3, \ldots \tag{2.56}$$

As x_- increases the first mass decreases less rapidly than the other levels (see Fig.2.7).

The GRS Region

For $x_- \gg e^{2x}$ the first mass is no longer special and scales with respect to *both* x and x_- in the same way as the remaining tower:

$$m_n = n\pi k\frac{e^{-x}}{x_-} \qquad n = 1, 2, 3, \ldots \tag{2.57}$$

The mass splittings Δm tend to zero as $x_- \to \infty$ and we obtain the GRS continuum of states (see Fig.2.7).

2.2 Multigravity

Armed with the details how the spectrum smoothly changes between the $''+-+''$ bigravity model ($x_- = 0$) and the GRS model ($x_- \to \infty$), we can now discuss the possibilities for modifying gravity at large distances. The couplings of the KK states with matter on one of the scale equivalent positive tension branes are readily calculated by (2.29):

$$a_n = \frac{1}{2\sqrt{2}}\left[\frac{g(0)}{M}\right]^{3/2}\hat{\Psi}^{(n)}(0) \tag{2.58}$$

Bigravity Region

As $x_- \to 0$, we are in the bigravity region which has been extensively explored in the previous section.

The GRS Region

In the GRS limit, $x_- \gg e^{2x}$, we should reproduce the "resonance"-like behaviour of the coupling in the GRS model. In the following we shall see that indeed this is the case and we will calculate the first order correction to the GRS potential for the case that x_- is large but finite.

For the rest of the section we split the wavefunction (2.51) in two parts, namely the normalization N_n and the unnormalized wavefunction $\psi^{(n)}(z)$, *i.e.* $\hat{\Psi}^{(n)}(z) = N_n\psi^{(n)}(z)$. The former is as usual chosen so that we get a canonically normalized Pauli-Fierz Lagrangian for the four dimensional KK modes $\bar{h}^{(n)}_{\mu\nu}$ (see (2.27)) and is given by:

$$N_n^2 = \frac{1/2}{2\displaystyle\int_0^{z_1} dz\left[\psi^{(n)}(z)\right]^2 + \int_{z_1}^{z_2} dz\left[\psi^{(n)}(z)\right]^2} \tag{2.59}$$

The value of $\psi^{(n)}(z)$ on the left positive tension brane is, for $m_n \ll k$:

$$\psi^2_{(n)}(0) = \frac{16k^3}{\pi^2 m_n^4} \tag{2.60}$$

It is convenient to split the gravitational potential given by the relation (2.32) into two parts:

$$V(r) = -\frac{1}{8M^3}\sum_{n=1}^{N_{x_-}-1}\frac{e^{-m_n r}}{r}N_n^2\psi^2_{(n)}(0) - \frac{1}{8M^3}\sum_{n=N_{x_-}}^{N_\Lambda}\frac{e^{-m_n r}}{r}N_n^2\psi^2_{(n)}(0) \tag{2.61}$$

where we denote by N_{x_-} the level after which the coupling of the KK states becomes linear to their mass. It corresponds to a mass scale about $m_s = ke^{-x}$. As we shall see this separation is useful because the first N_{x_-} states give rise to the long distance gravitational potential V_{ld} while the remaining ones will only contribute to the short distance corrections ΔV_{sd}.

- **Short Distance Corrections**

We first consider the second term. The normalization constant in this region is computed by considering the asymptotic expansions of the Bessel functions with argument $\frac{g(z_1)m_n}{k}$. It is calculated to be:

$$N_n^2 = \frac{\pi^3 m_n^5}{32k^3 g(z_1) x_-} \left[\frac{1}{1+\frac{2}{x_-}}\right] \tag{2.62}$$

If we combine the above normalization constant with the unnormalized wavefunction (2.60), we find:

$$\Delta V_{sd}(r) = -\frac{1}{32\pi M^3} \sum_{n=N_{x_-}}^{N_\Lambda} \frac{k\pi}{x_- e^x} \frac{m_n}{2k} \frac{e^{-m_n r}}{r} \left[\frac{1}{1+\frac{2}{x_-}}\right] \tag{2.63}$$

Since we are taking $x_- \gg e^{2x}$, the spectrum tends to continuum, *i.e.* $N_n \to N(m)$, $\psi_{(n)}(0) \to \psi(m)$, and the sum turns to an integral where the first factor in (2.63) is the integration measure, *i.e.* $\sum \frac{k\pi}{x_- e^x} = \sum \Delta m \to \int dm$ from (2.57). Moreover, we can extend the integration to infinity because the integral saturates rather quickly. Finally, we expand the fraction involving x_- keeping the first term in the power series to obtain the potential:

$$\Delta V_{sd}(r) = -\frac{1}{32\pi M^3} \int_{m_s}^{\infty} dm \, \frac{e^{-mr}}{r} \, \frac{m}{2k} \left(1 - \frac{2}{x_-}\right) \tag{2.64}$$

The integral is easily calculated and the potential reads:

$$\Delta V_{sd}(r) = -\frac{G_N}{2r} \, \frac{1+m_s r}{(kr)^2} \, (1 - \frac{2}{x_-}) \, e^{-m_s r} \tag{2.65}$$

where we identified $G_N \equiv \frac{k}{32\pi M^3}$ for reasons to be seen later. The second part of the above potential is the first correction coming from the fact that x_- is finite. Obviously this correction vanishes when $x_- \to \infty$. Note that the above potential gives corrections to the Newton's law only at distances comparable to the Planck length scale.

- **Multigravity: 4D and 5D gravity**

We turn now to the more interesting first summation in (2.61) in order to show that the coupling indeed has the "resonance"-like behaviour for $\Delta m \to 0$ responsible for four dimensional Newtonian gravity at intermediate distances and the five dimensional gravitational law for cosmological distances. This summation includes the KK states from the graviton zero mode up to the N_{x_-}-th level. The normalization constant in this region is computed by considering the series expansion of all the Bessel functions involved. It is calculated to be:

$$N_n^2 = \frac{\pi^2 m_n^4}{4g(z_1)^4 x_-} \left[\frac{1}{m_n^2 + \frac{\Gamma^2}{4} + \frac{8k^2}{g(z_1)^4 x_-}}\right] \tag{2.66}$$

where $\Gamma = 4ke^{-3x}$. If we combine the above normalization constant with the unnormalized wavefunction (2.60), we find that the long distance gravitational potential is:

$$V_{ld}(r) = -\frac{1}{32\pi M^3} \sum_{n=0}^{N_{x_-}} \frac{\pi k}{x_- e^x} \frac{4k^2}{\pi g(z_1)^3} \frac{e^{-m_n r}}{r} \left[\frac{1}{m_n^2 + \frac{\Gamma^2}{4} + \frac{8k^2}{g(z_1)^4 x_-}}\right] \tag{2.67}$$

Again, since we are taking $x_- \gg e^{2x}$, the above sum will turn to an integral with $\sum \Delta m \to \int dm$. Moreover, we can safely extend the integration to infinity since the integral saturates very fast for $m \lesssim \Gamma/4 \equiv r_c^{-1} \ll ke^x$. If we also expand the fraction in brackets keeping the first term in the power series, we find the potential:

$$\begin{aligned} V_{ld}(r) = & - \frac{1}{32\pi M^3} \int_0^\infty dm \, \frac{4k^2}{\pi g(z_1)^3} \frac{e^{-mr}}{r} \frac{1}{m^2 + \frac{\Gamma^2}{4}} \\ & + \frac{1}{32\pi M^3} \int_0^\infty dm \, \frac{32k^4}{x_- \pi g(z_1)^7} \frac{e^{-mr}}{r} \frac{1}{(m^2 + \frac{\Gamma^2}{4})^2} \end{aligned} \tag{2.68}$$

The first part is the same as in the GRS model potential, whereas the second one is the first correction that comes from the fact that x_- is still finite though very large. Note that the width of the "resonance" scales like e^{-3x}, something that is compatible with the scaling law of the masses ($m_n = n\pi k \frac{e^{-x}}{x_-}$), since we are working at the region where $x_- \gg e^{2x}$, *i.e.* $m_n \ll n\pi k e^{-3x}$. The above integrals can be easily calculated in two interesting limits:

For $k^{-1} \ll r \ll r_c$ the potential is given approximately by:

$$V_{ld}(r \ll r_c) = -\frac{G_N}{r} \left(1 - \frac{e^{2x}}{x_-}\right) \tag{2.69}$$



Figure 2.8: The behaviour of the coupling, $a(m)$, in the limit of $x_- \gg e^{2x}$. Three regions of interest are indicated. The region $m > m_s$ gives rise to short distance corrections. The $m_1 \ll m \ll m_c$ region gives rise to 4D gravity at intermediate distances and 5D gravity at ultra large distances. For distances $r \gg m_1^{-1}$, the zero mode gives the dominant contribution and thus we return to 4D gravity.

where we have identified $G_N \equiv \frac{k}{32\pi M^3}$ to obtain the normal four dimensional Newtonian potential. Note that since $x_- \gg e^{2x}$, the $1/x_-$ term is indeed a small correction.

In the other limit, $r \gg r_c$, the integrand is only significant for values of m for which the m^2 term in the denominator of the Breit-Wigner can be dropped and the potential becomes five dimensional:

$$V_{ld}(r \gg r_c) = -\frac{G_N r_c}{\pi r^2}\,(1 - \frac{2e^{2x}}{x_-}) \tag{2.70}$$

The fact that Newtonian gravity has been tested close to the present horizon size require that for $k \sim M_{\rm Pl}$ we should have $x \gtrsim$ 45-50. Finally, we note that if we take the $x_- \to \infty$ we recover the GRS result:

$$\lim_{x_- \to \infty} V_{+--+}(r, x_-) = V_{GRS}(r) \tag{2.71}$$

- **Back to 4D gravity**

As we have just seen, probing larger distances than r_c, the four dimensional gravitational potential changes to a five dimensional one. This is the most significant characteristic of the GRS model. In the case that x_- is large compared to e^{2x} but still finite, there is another distinct region of interest, namely $r \gg m_1^{-1}$. This follows from the fact that, in this limit, the spectrum is still discrete. For distances larger than of the order of the corresponding wavelength of the first KK mode, the contribution to gravity from the KK tower is suppressed and thus the zero mode gives the dominant contribution, leading to the four dimensional Newtonian potential again. In this case the strength of the gravitational interaction is a small fraction of the strength of the intermediate scale four dimensional gravity. More precisely, the contribution of the massless graviton is $1/x_-$ suppressed and thus vanishes when $x_- \to \infty$, something that is expected since in this limit there is no nomalizable zero mode. The gravitational potential in this case is:

$$V_{4D}(r) = -\frac{1}{32\pi M^3}\,\frac{1}{r}\;N_0^2\psi_{(0)}^2(0) = -\frac{G_N}{r}\;\frac{e^{2x}}{x_-} \tag{2.72}$$

Obviously this four dimensional region disappears at the limit $x_- \to \infty$ since the spectrum becomes continuum and thus the five dimensional gravity "window" extents to infinity. We should note finally that for the values of x that we consider here this final modification of gravity occurs at distances well above the present horizon.

3 The Crystal Universe Model

The last model we consider is the crystal universe introduced in [54] (see also [55,56]). It consists of an infinite array of parallel three-branes in five dimensional AdS space with cosmological constant Λ. For simplicity we assume that the branes are equidistant with distance, l, between two successive branes. Needless to say, in this case all the positive tension branes are scale equivalent. The metric ansatz that respects four dimensional Poincaré invariance is again given by (2.3) where the $\sigma(y)$ function is constrained by the the Einstein equations to have the sawtooth form:

$$\sigma(y) = k\sum_{j=-\infty}^{+\infty}(y-2jl)\left[2\theta(y-2jl)-\theta(y-(2j-1)l)-\theta(y-(2j+1)l)\right] \tag{2.73}$$

The tensions of successive branes are required to be opposite and equal to $\pm\Lambda/k$, where $k=\sqrt{\frac{-\Lambda}{24M^3}}$ is a measure of the curvature of the bulk and M the five dimensional fundamental scale.

We consider the general fluctuations around the vacuum ansatz of the form (2.10) and decompose as usual the perturbation tensor $h_{\mu\nu}(x,y)$ as:

$$h_{\mu\nu}(x,y) = \int dm\, h_{\mu\nu}(m,x)\Psi(m,y) \tag{2.74}$$



Figure 2.9: The Crystal Universe made up of an infinite array of "+" and "−" branes with lattice spacing l and bulk curvature k.

where $\Psi(m, y)$ is a complex function that arises from the linear combination of the independent graviton polarizations. This leads to the Schrödinger equation (2.13) for the wavefunction $\hat{\Psi}(m, z) \equiv \Psi(m, y)e^{\sigma/2}$ with potential:

$$V(z) = \frac{15k^2}{8[g(z)]^2} - \frac{3k}{2g(z)} \sum_{j=-\infty}^{+\infty} (-)^j \delta(z - z_i) \tag{2.75}$$

The z coordinates have been defined as usual to be the ones that make the background metric conformally flat. In these coordinates the branes sit at the points $z_j = j\frac{e^{kl}-1}{k} \equiv jz_l$ and the function $g(z)$ is:

$$g(z) = 1 + k \sum_{j=-\infty}^{+\infty} (z - z_{2j}) \left[2\theta(z - z_{2j}) - \theta(z - z_{2j-1}) - \theta(z - z_{2j+1})\right] \tag{2.76}$$

The solution of the Schrödinger equation for the wavefunctions for two adjacent cells is given [54] in terms of Hankel functions:

$$\hat{\Psi}(m, z) = N_m \sqrt{\frac{g(z)}{k}} \left[\left\{ \begin{array}{c} 1 \\ B \\ e^{2iqz_l} \\ e^{2iqz_l} B \end{array} \right\} H_2^+ \left(\frac{m}{k} g(z)\right) + \left\{ \begin{array}{c} A \\ C \\ e^{2iqz_l} A \\ e^{2iqz_l} C \end{array} \right\} H_2^- \left(\frac{m}{k} g(z)\right) \right] \tag{2.77}$$

in the regions $z \in [0, z_l]$, $z \in [z_l, 2z_l]$, $z \in [2z_l, 3z_l]$ and $z \in [3z_l, 4z_l]$ respectively. In the above expression N_m is an overall normalization constant, q is the Bloch wave quasi-momentum and the constants A,B,C are given by:

$$\begin{aligned} A &= \frac{(h_1^+h_2^- + h_1^-h_2^+)\hat{h}_2^+ - (h_1^-h_2^+ - h_1^+h_2^-)e^{2iqz_l}\hat{h}_2^+ - 2h_1^+h_2^+\hat{h}_2^-}{(h_1^+h_2^- + h_1^-h_2^+)\hat{h}_2^- + (h_1^-h_2^+ - h_1^+h_2^-)e^{2iqz_l}\hat{h}_2^- - 2h_1^-h_2^-\hat{h}_2^+} && (2.78) \\ B &= \frac{h_1^+h_2^- + h_1^-h_2^+ + 2h_1^-h_2^-A}{h_1^-h_2^+ - h_1^+h_2^-} && (2.79) \\ C &= \frac{2h_1^+h_2^+ + (h_1^+h_2^- + h_1^-h_2^+)A}{h_1^+h_2^- - h_1^-h_2^+} && (2.80) \end{aligned}$$

where $h_n^\pm \equiv H_n^\pm\left(\frac{m}{k}g(z_l)\right)$ and $\hat{h}_n^\pm \equiv H_n^\pm\left(\frac{m}{k}\right)$. The above coefficients of the Hankel functions were determined by the boundary conditions, *i.e.* continuity of the wavefunction at z_l, $2z_l$, $3z_l$, discontinuity of its first derivative at z_l, $3z_l$ and the Bloch wave conditions relating the wavefunction at the edges of each cell[2]. The last remaining boundary condition of the discontinuity of the first derivative of the wavefunction at $2z_l$ gives us the band equation connecting q and m. We disagree with the relation given in [54] and instead we find:

$$\cos(2qz_l) = \frac{(j_2y_1 + j_1y_2)(\hat{j}_2\hat{y}_1 + \hat{j}_1\hat{y}_2) - 2\hat{j}_1\hat{j}_2y_1y_2 - 2j_1j_2\hat{y}_1\hat{y}_2}{(j_2y_1 - j_1y_2)(\hat{j}_2\hat{y}_1 - \hat{j}_1\hat{y}_2)} \equiv f(m) \qquad (2.81)$$

where again $j_n \equiv J_n\left(\frac{m}{k}g(z_l)\right)$, $y_n \equiv Y_n\left(\frac{m}{k}g(z_l)\right)$ and $\hat{j}_n \equiv J_n\left(\frac{m}{k}\right)$, $\hat{y}_n \equiv Y_n\left(\frac{m}{k}\right)$. This dispersion relation *always* gives a band at zero as is to be expected intuitively. Defining the parameter $x \equiv kl$, we see that for $x \gtrsim 10$ we may reliably approximate the width of the zero mode band by:

$$\Gamma_0 = 2\sqrt{2}ke^{-2x} \qquad (2.82)$$

while the separation of the zeroth and the first band is

$$\Delta\Gamma_1 = \xi_1 ke^{-x} \qquad (2.83)$$

with ξ_1 the 1st root of $J_1(x)$. This characteristic behaviour of the above widths with respect to x implies that we can have a viable multigravity scenario if $\Gamma_0^{-1} \gtrsim 10^{26}$cm and at the same time $\Delta\Gamma_1^{-1} \lesssim 0.2$mm. This can happen for $k \sim M_{\rm Pl}$ and $x \approx 68$.

The band structure has the following form. The first bands have very narrow widths Γ_i with spacing $\Delta\Gamma_i$ between them (*i.e.* between the i-th and the $(i-1)$-th band), which are approximately:

---

[2] The Bloch wave condition reads $\hat{\Psi}(z_i + 2z_l) = e^{2iqz_l}\hat{\Psi}(z_i)$.

$$\Gamma_i = \epsilon_i \xi_i^2 k e^{-3x} \tag{2.84}$$

$$\Delta\Gamma_i = (\xi_i - \xi_{i-1}) k e^{-x} \tag{2.85}$$

where ξ_{2n+1} is the $(n+1)$-th root of $J_1(x)$, ξ_{2n} is the n-th root of $J_2(x)$, and $\epsilon_{2n+1} = \frac{\pi}{4}\frac{Y_1(\xi_{2n+1})}{J_2(\xi_{2n+1})}$, $\epsilon_{2n} = \frac{\pi}{4}\frac{Y_2(\xi_{2n})}{J_1(\xi_{2n})}$. As we move on to higher bands their width is increasing and the spacing between them is decreasing. In the limit $m \to \infty$, the spacing disappears, *i.e.* $\Delta\Gamma_i \to 0$, while their width tends to $\Gamma_i \to \frac{\pi}{2} k e^{-x}$. In this limit the function $f(m)$ in (2.81) tends to $f(m) = \cos(2 z_l m)$ and we have no forbidden zones.

In order to find the gravitational potential we should at first find how the KK modes couple to matter and determine their spectral density. Instead of working with wavefunctions in an infinite extra dimension with continuous normalization, it is more convenient effectively to compactify the system to a circle of length $2\mathcal{N} l$, by assuming that we have a finite crystal of $\mathcal{N}$ cells $[2jl, 2(j+1)l]$. We shall find the gravitational potential is independent of $\mathcal{N}$, so we will be able to take the limit $\mathcal{N} \to \infty$ and decompactify the geometry.

In this regularization scheme we can readily see that the ever-present zero mode, corresponding to the four dimensional graviton, has wavefunction:

$$\hat{\Psi}(0, z) = \frac{\sqrt{k/\mathcal{N}}}{[g(z)]^{3/2}} \tag{2.86}$$

and its coupling to matter on the central positive brane is:

$$a(0) = \frac{1}{\sqrt{\mathcal{N}}} \frac{\sqrt{k}}{2\sqrt{2M^3}} \equiv \frac{1}{\sqrt{\mathcal{N}}} \frac{1}{2\sqrt{2}M_*} \tag{2.87}$$

The wavefunction (2.86) is a smooth limit of (2.77) for $m \to 0$. Indeed, for large x we find the normalization constant in (2.77) in the region of the zero mode band is approximately:

$$N_m = \frac{\sqrt{2}}{z_l^2 m} \frac{1}{\sqrt{\mathcal{N}}} \tag{2.88}$$

The approximation of the Hankel functions with their first term of their power series proves our previous claim. Furthermore, we can see that the KK states in the whole zero mode band for large x will couple to matter in the central positive brane as:

$$a(m) = \frac{1}{\sqrt{\mathcal{N}}}\frac{1}{2\sqrt{2}M_*}e^{i\phi(m)} \tag{2.89}$$

with $\phi(m)$ an unimportant m-dependent phase that does not appear in observable quantities.

In the limit $m \to \infty$ the Hankel functions can be approximated by their asymptotic form. The wavefunction is then almost constant in z-space and the KK states couple to matter on the central positive brane as:

$$a(m) = \frac{1}{\sqrt{\mathcal{N}}}\frac{1}{4e^{x/2}M_*}e^{i\phi(m)} \tag{2.90}$$

where again $\phi(m)$ is an unimportant m-dependent phase. The couplings in the low lying bands interpolate between the above limiting cases.

The standard procedure to calculate the spectral density is at first to find the number of states with masses smaller than m and then differentiate with respect to m. In our regularization prescription it is straightforward to count the states in the quasi-momentum space with masses lighter that m. The latter is simply:

$$N_{tot}(m) = \frac{\mathcal{N} z_l}{\pi} q \tag{2.91}$$

so the spectral density is its derivative:

$$\rho(m) = \frac{dN_{tot}(m)}{dm} = \frac{\mathcal{N} z_l}{\pi}\frac{dq}{dm} = (\mp)\frac{\mathcal{N}}{2\pi}\frac{1}{\sqrt{1-f(m)^2}}\frac{df(m)}{dm} \tag{2.92}$$

where the plus and minus signs will succeed each other for neighbouring bands, starting with minus for the zero mode band. For the zero band we can have a reliable approximation for large x of the above function:

$$\rho(m) = \mathcal{N}\frac{1}{\pi\Gamma_0}\frac{1}{\sqrt{1-(m/\Gamma_0)^2}} \tag{2.93}$$

Thus, the spectral density diverges as $m \to \Gamma_0$ but the divergence is integrable and does not cause any problem to the following calculations. Actually, the spectral density diverges at the edges of every band since at these points $\sqrt{1-f(m)^2} \to 0$ while $\frac{df(m)}{dm} \neq 0$. The only point that this does not happen is at $m = 0$ where $\frac{df(m)}{dm} \to 0$ in such a way that the result is finite.

In the limit $m \to \infty$ the bands disappear and the spectrum has constant spectral



Figure 2.10: Spectral density for the first three bands in the crystalline model.

density:

$$\rho(m) = \mathcal{N}\frac{e^x}{k\pi} \tag{2.94}$$

The gravitational potential, taking into account the Yukawa suppressions of the KK states, is simply:

$$V(r) = -\int_0^\infty dm \frac{|a(m)|^2}{4\pi}\rho(m)\frac{e^{-mr}}{r} \tag{2.95}$$

For distances $r \gg \Delta\Gamma_1^{-1}$ we have the effective potential:

$$\begin{aligned} V(r) &= -\frac{1}{32\pi M_*^2}\frac{1}{r}\int_0^{\Gamma_0} dm \frac{\rho(m)}{\mathcal{N}} e^{-mr} = -\frac{1}{32\pi^2 M_*^2}\frac{1}{r}\int_0^1 d\xi \frac{e^{-(\Gamma_0 r)\xi}}{\sqrt{1-\xi^2}} \\ &= -\frac{1}{64\pi M_*^2}\frac{1}{r}[I_0(\Gamma_0 r) - L_0(\Gamma_0 r)] \end{aligned} \tag{2.96}$$

where I_0 is the 0-th modified Bessel function and L_0 is the 0-th modified Struve function. The gravitational potential is $\mathcal{N}$-independent as expected. For distances $\Delta\Gamma_1^{-1} \ll r \ll \Gamma_0^{-1}$ the above functions tend to $I_0(\Gamma_0 r) \to 1$ and $L_0(\Gamma_0 r) \to 0$, so we recover the four dimensional Newton law with $G_N = \frac{1}{64\pi M_*^2}$:

$$V(r \ll \Gamma_0^{-1}) = -\frac{G_N}{r} \tag{2.97}$$

On the other hand, for distances $r \gg \Gamma_0^{-1}$ the asymptotic expansion of the difference of the modified Bessel and Struve functions gives a five dimensional Newton's law:

$$V(r \gg \Gamma_0^{-1}) = -\frac{2G_N}{\pi\Gamma_0}\frac{1}{r^2} \tag{2.98}$$

In case we take the crystal to be finite, there will appear a region for $r \gg m_1^{-1}$ where gravity will turn again to four dimensional. However, as explained in the previous section, this region is well above the universe horizon and thus of no phenomenological interest.

Finally, for distances $r \lesssim \Delta\Gamma_1^{-1}$ we will start to feel the short distance Yukawa type modifications to gravity due to the presence of the bands above the zeroth one. As in the zero hierarchy bigravity and $'' + - - +''$ multigravity model we expect these corrections to be important at scales of the order of the Planck length. In summary, the physics of the crystalline model is very similar to the GRS model as far as modifications of gravity due to tensor modes is concerned. In both cases we have a crossover from four dimensional to five dimensional gravity at cosmological scales.

4 Propagator structure

We have shown that in models in which negative tension branes are sandwiched between positive tension branes there are anomalously light KK states that can lead to modifications of gravity. So far we have implicitly assumed that the gravitational force generated by a massive graviton is identical to that generated by a massless one but this is not, in general, the case because the massive graviton has additional degrees of freedom which do not decouple in the massless limit. As a result the interaction generated by a massive graviton violates the normal four dimensional relation between the gravitational interactions of matter and light in Einsteinian gravity [29,30] . This may be seen explicitly from the form of the massive and massless graviton propagators.

In more detail, the form of the massless graviton propagator in flat spacetime (in momentum space) has the form:

$$G^{\mu\nu;\alpha\beta} = \frac{1}{2}\,\frac{\left(\eta^{\mu\alpha}\eta^{\nu\beta} + \eta^{\nu\alpha}\eta^{\mu\beta}\right) - \eta^{\mu\nu}\eta^{\alpha\beta}}{p^2} + \cdots \tag{2.99}$$

where we have omitted terms that do not contribute when contracted with a conserved energy momentum tensor $T_{\mu\nu}$.

On the other hand, the four-dimensional massive graviton propagator (in momentum space) has the form:

$$G^{\mu\nu;\alpha\beta} = \frac{1}{2}\,\frac{\left(\eta^{\mu\alpha}\eta^{\nu\beta} + \eta^{\nu\alpha}\eta^{\mu\beta}\right) - \frac{2}{3}\,\eta^{\mu\nu}\eta^{\alpha\beta}}{p^2 + m^2} + \cdots \tag{2.100}$$

The difference between the two propagators

$$\delta G^{\mu\nu,\alpha\beta} = \frac{1}{6}\,\frac{g^{\mu\nu}g^{\alpha\beta}}{p^2 + m^2} + \cdots \tag{2.101}$$

is due to the additional helicity components needed for a massive graviton. There are three additional components needed. Two, corresponding to the graviphotons, decouple at low energies as they are derivatively coupled to the conserved energy-momentum tensor. The third, corresponding to a scalar component, does not decouple and is responsible for δG. The contribution of δG to the one-graviton exchange amplitude for any two four dimensional sources $T^1_{\mu\nu}$ and $T^2_{\alpha\beta}$ vanishes for light for which $T = T^\mu_\mu = 0$ but not for matter because $T = T^0_0 \neq 0$, hence the normal four dimensional gravity relation between the interactions of matter and light is violated as was first pointed out by [24]. For example, if gravity is due to the exchange of a massive spin 2 particle and one insists in the validity of Newton's law for static sources, then the deflection angle of light would be 25% smaller than if it corresponds to the exchange of the massless graviton. The fact that the bending of the light by the sun agrees with the prediction of Einstein's theory to 0.02% accuracy, rules out the possibility that gravity is due to massive graviton exchange irrespective of how small its mass is. Of course there remains the possibility that a small fraction of the gravitational interactions are associated with a massive graviton component in the presence of a dominant massless graviton component. This can be realized by having an ultralight spin-2 particle with a very small coupling compared to graviton's. Such an example could be the massless - massive graviton system in the asymmetric bigravity model case.

However, in our discussion so far we have omitted an important gravitational mode, the moduli (radions) corresponding to moving negative tension branes. These radions as we shall see in the following chapter cancel the troublesome additional polarization states of the massive gravitons [37, 36, 32, 33] . This restores the phenomenologically favoured propagator structure of the above models at intermediate distances. However, these fields must have unusual properties because it is known that a normal scalar contributes with the same sign as δG and therefore cannot cancel it. An explicit computation presented in the following chapter reveals that it has a negative kinetic energy term leading to a contribution which cancels the troublesome δG contribution. As a result the theory indeed gives rise to normal four dimensional gravity at intermediate distances, but at large distances the ghost radions dominate giving in all cases scalar antigravity [37]. The presence of the ghost radion field is worrying for the consistency of the theory. We will elaborate on this issue in the following chapter.

Chapter 3

Radions and dilaton

In this chapter we are interested in describing the dynamics of the scalar gravitational perturbations for a system with branes in a warped background in five dimensions. These excitations describe the effect of the fluctuation of the size of the extra dimension and/or of the relative positions of the branes. We will distinguish these two kinds of modes by calling the former dilaton [34] and the latter radions [34,32,33]. An orbifold symmetry Z_2 acting on the extra dimensional coordinate as $y \rightarrow -y$ is also imposed. When the topology of the extra dimension is S^1, the compact case, the Z_2 action has two fixed points $y = 0,\ L$. If branes are sitting on the orbifold fixed points, the corresponding radions are frozen. For example in a system of three three-branes where two of them are sitting on fixed points at $y = 0,\ L$, we are left with just one radion field corresponding to the fluctuation of the position of the third brane and the dilaton corresponding to the fluctuation of the size of the orbifold.

For the sake of simplicity we will only consider a three flat three-branes system embedded in AdS_5 with in general different cosmological constants between the branes. This will be sufficient to clarify the role of the radion and dilaton in the multibrane models. We will calculate the full gravitational effective action for this general model, compute the kinetic energy of the moduli and study the nature of gravity on the branes. As we have already noted the radion excitations play an important role in the context of multigravity models. They restore the correct propagator structure for the massive graviton excitations. However, the radions corresponding to negative tension branes are physical ghosts and therefore disastrous for the theory.

Afterwards, we will concentrate on the dilaton of a five dimensional model with identical four geometries and calculate its mass for bulk dynamics involving arbitrary minimally coupled scalar fields. A useful tool for this calculation is the sum rules derived by Gibbons, Kallosh and Linde [41] which we briefly review. We find a generic formula relating the dilaton mass with the sum of the tensions of the branes and the curvature of the four-geometry. Demanding that this mass is not tachyonic we can derive the necessary condition for stabilizing the overall size of the system.



Figure 3.1: General three three-brane model on an orbifold with different cosmological constants between the branes.

1 The general three three-brane system

We will consider a three three-brane model on an S^1/Z_2 orbifold. Two of the branes sit on the orbifold fixed points $y = y_0 = 0$, $y = y_2 = L$ respectively. A third brane is sandwiched in between at position $y = y_1 = r$ as in Fig.3.1. In each region between the branes the space is AdS_5 and in general the various AdS_5 regions have different cosmological constants Λ_1, Λ_2. The action describing the above system is:

$$\mathcal{S}_G = \int d^4x dy \sqrt{-G} \left[2M^3 R - \Lambda(y) - \sum_i V_i \, \delta(y - y_i) \frac{\sqrt{-\hat{G}^{(i)}}}{\sqrt{-G}} \right] \tag{3.1}$$

where M is the five dimensional Planck mass, V_i the tensions of the gravitating branes and $\hat{G}^{(i)}_{\mu\nu}$ the induced metric on the branes. The orbifold symmetry $y \to -y$ is imposed.

We seek a background static solution of Einstein equations for the following four dimensional Poincaré invariant metric ansatz:

$$ds^2 = a^2(y) \, \eta_{\mu\nu} dx^\mu dx^\nu + dy^2 \tag{3.2}$$

The solution for the warp factor has the usual exponential form:

$$a(y) = \begin{cases} e^{-k_1 y} & , 0 < y < r \\ e^{-k_2 y + r(k_2 - k_1)} & , r < y < L \end{cases} \tag{3.3}$$

where k_1 and k_2 are the curvatures of the bulk in the two regions and are related to the bulk cosmological constants as:

$$k_1^2 = -\frac{\Lambda_1}{24M^3}\,, \qquad k_2^2 = -\frac{\Lambda_2}{24M^3} \tag{3.4}$$

Moreover, the Einstein equations impose the following fine tuning between the brane tensions and the bulk cosmological constants:

$$V_0 = 24M^3\,k_1\,, \qquad V_1 = 24M^3\frac{(k_2-k_1)}{2}\,, \qquad V_2 = -24M^3\,k_2 \tag{3.5}$$

It is straightforward to recover some special models that have been considered in the literature. The RS1 model is obtained for $k_1 = k_2$ where the intermediate brane is absent (zero tension). For $k_1 > 0$ and $k_2 = -k_1$ we get the $''+-+''$ bigravity model [21, 53]. For $k_1 > 0$ and $k_2 > k_1$ we get the $''++-''$ brane model considered in [26]. In the decompactification limit where $L \to \infty$ we get also two interesting models: For $k_1 > 0$ and $k_2 = 0$ we obtain the GRS model [22] and for $k_1 > 0$ and $k_2 = 0 > k_1$ the non-zero tension version [32] of the model considered in [57].

2 Effective action

Our purpose is to study fluctuations of the background (3.2). The first important observation is that there exists a generalization of Gaussian normal coordinates such that in the perturbed geometry the $g_{\mu y}$ components of the metric are zero and at the same time the embedding of branes is still described by $y = 0$, $y = r$ and $y = L$ (see for instance the Appendix of [32]). In that case the metric takes the form:

$$ds^2 = g_{\mu\nu}(x,y)\,dx^\mu dx^\nu + g_{yy}(x,y)\,dy^2 \quad . \tag{3.6}$$

When analyzed from a four dimensional point of view, in each region, perturbations are of of three types.

- **Spin two**:

  Tensor-like perturbation $\tilde{h}_{\mu\nu}(x,y)$ corresponding to massive (and massless) four dimensional gravitons[1]:

$$ds^2 = a^2(y)\left[\eta_{\mu\nu} + \tilde{h}_{\mu\nu}(x,y)\right]\,dx^\mu dx^\nu + dy^2 \tag{3.7}$$

---

[1] We denote $\tilde{h}_{\mu\nu}$ the conformal perturbation in contrast to the previous chapter graviton perturbation $h_{\mu\nu}$. Their relation is $h_{\mu\nu} = a^2\tilde{h}_{\mu\nu}$.

- **Spin zero: Dilaton**

  Scalar perturbation $f_1(x)$ corresponding to the fluctuation of the size of the orbifold [34]:

$$ds^2 = a^2(y)\,[1 + Q(y)f_1(x)]\,\eta_{\mu\nu}\,dx^\mu dx^\nu + [1 + q(y)f_1(x)]\,dy^2 \tag{3.8}$$

- **Spin zero: Radion**

  Scalar perturbation $f_2(x)$ corresponding to a fluctuating distance of the intermediate brane [32]:

$$ds^2 = a^2(y)\,[(1 + B(y)f_2(x))\,\eta_{\mu\nu} + 2\epsilon(y)\,\partial_\mu\partial_\nu f_2(x)]\,dx^\mu dx^\nu + [1 + A(y)\,f_2(x)]\,dy^2 \tag{3.9}$$

In the above decomposition we should note that we have kept only the zero modes of the radion and the dilaton. Their massive KK-states are "eaten" by the graviton's corresponding modes in order to give them mass, in a geometrical realization of the Higgs phenomenon. The function $\epsilon(y)$ is needed in order that the metric satisfies the Israel junction conditions in the presence of moving branes. As it will be shown later, the values of $\partial_y\epsilon$ at $y = 0$, r, L are gauge invariant. When $k_1 \neq k_2$ a non-trivial ϵ is required.

The generic perturbation can be written as:

$$\begin{aligned} ds^2 =& a^2(y)\left\{[1 + \varphi_1(x,y)]\,\eta_{\mu\nu} + 2\epsilon(y)\,\partial_\mu\partial_\nu f_2(x) + \tilde{h}_{\mu\nu}(x,y)\right\} dx^\mu dx^\nu \\ &+ [1 + \varphi_2(x,y)]\,dy^2 \end{aligned} \tag{3.10}$$

where

$$\begin{aligned} \varphi_1(x,y) &= Q(y)f_1(x) + B(y)f_2(x) \\ \varphi_2(x,y) &= q(y)f_1(x) + A(y)f_2(x) \end{aligned} \tag{3.11}$$

Given the expression (3.3) for a, the Israel junctions conditions (continuity of metric components and jump conditions) at $y = 0, r, L$, simply require that A, B, $\partial_y\epsilon$, Q, q are continuous there [32]. The four dimensional effective action $\mathcal{S}_{eff}$ for the various modes is obtained inserting the ansatz (3.11) in the action (3.1) and integrating out y. So far the functions A, B, ϵ, Q, q have not been specified, however imposing that $\mathcal{S}_{eff}$ contains no mixing terms among $\tilde{h}$ and f_i one determines Q, q and finds A in terms of B (see Appendix A). The latter satisfies:

$$\frac{d}{dy}\left(Ba^2\right)+2a^{-1}\frac{da}{dy}\frac{d}{dy}\left(a^4\partial_y\epsilon\right)=0 \tag{3.12}$$

$$\int_{-L}^{L}dy\, a\left(\frac{da}{dy}\right)^{-1}\frac{dB}{dy}=0 \tag{3.13}$$

The condition that $\tilde{h}$ and f_i do not mix between them allows us to give a clear particle interpretation for the quanta of $\tilde{h}$ and f_i. However, despite the fact that we have chosen that gauge, the results that we obtain are bound to be the same as in the general case where there would be quadratic mixing between the modes, because of the five dimensional gauge invariance of the theory. As a consequence of the no-mixing conditions the linearized Einstein equations for (3.11) will consist in a set independent equations for the graviton and the scalars.

The effective Lagrangian reads (see Appendix A):

$$\begin{aligned}\mathcal{S}_{eff}&=\int d^4x\,\mathcal{L}_{eff}=\int d^4x\,\left(\mathcal{L}_{Grav}+\mathcal{L}_{Scal}\right)\\ \mathcal{L}_{eff}&=2M^3\int_{-L}^{L}dy\left\{a^2\mathcal{L}_{PF}^{(0)}(\tilde{h})+\frac{a^4}{4}\left[(\partial_y\tilde{h})^2-\partial_y\tilde{h}_{\mu\nu}\,\partial_y\tilde{h}^{\mu\nu}\right]+\mathcal{L}_{Scal}\right\}\end{aligned} \tag{3.14}$$

with

$$\begin{gathered}\mathcal{L}_{Scal}=\mathcal{K}_1\,f_1\Box f_1+\mathcal{K}_2\,f_2\Box f_2\\ \mathcal{K}_1=2M^3\frac{3}{2}c^2\int_{-L}^{L}a^{-2}\,dy\,,\quad \mathcal{K}_2=-2M^3\,\frac{3}{4}\int_{-L}^{L}a\left(\frac{da}{dy}\right)^{-1}\frac{d}{dy}\left(B^2a^2\right)\,dy\end{gathered} \tag{3.15}$$

where c is an integration constant.

In (3.14), the spin-2 part, as expected, contains the four dimensional Pauli-Fierz Lagrangian $\mathcal{L}_{PF}^{(0)}(\tilde{h})$ for the gravitons without the mass term (see Appendix A), plus a mass term coming from the dimensional reduction. In the scalar part $\mathcal{L}_{Scal}$ the mass terms are zero since f_i are moduli fields. The indices in the above expressions are raised and lowered with $\eta_{\mu\nu}$. Notice that after the no-mix conditions are enforced, the effective Lagrangian contains the undetermined function $\epsilon(y)$.

The metric ansatz G_{MN} in (3.11) is related to a special coordinate choice, nevertheless, a residual gauge (coordinate) invariance is still present. Consider the class of infinitesimal coordinate transformations $x^M \to x'^M = x^M + \xi^M(x)$ such that the transformed metric ${G_{MN}}' = G_{MN} + \delta G_{MN}$ retains the original form (3.10) up to a redefinition of the functions q, Q, A, B, ϵ and the dilaton and the radion field. Consistency with the orbifold geometry and the requirement that the brane in $y = r$ is kept fixed by the diffeomorphism lead to $\xi^5(x,0) = \xi^5(x,r) = \xi^5(x,L) = 0$. From

$$\delta G_{MN} = -\xi^A\,\partial_A G_{MN} - \partial_M \xi^A\, G_{AN} - \partial_N \xi^A\, G_{MA} \tag{3.16}$$

one can show that ξ^M has to be of the form:

$$\begin{gathered}\xi^\mu(x,y) = \hat{\xi}^\mu(x) - W(y)\,\eta^{\mu\nu}\,\partial_\nu f_2(x)\,, \qquad \xi^5(x,y) = a^2 W'(y)\, f_2(x) \\ \text{with } \; W'(0) = W'(r) = W'(L) = 0\end{gathered} \tag{3.17}$$

The case $W = 0$ corresponds the familiar pure four dimensional diffeomorphisms, under which $\tilde{h}_{\mu\nu}$ transforms as spin two field, f_i as scalars and q, Q, A, B, ϵ are left unchanged. On the contrary the case $\hat{\xi}^\mu = 0$, $W \neq 0$ is relic of five dimensional diffeomorphisms and one can check that q, Q, A, B, ϵ are not invariant and in particular $\delta\epsilon = W$. As a result, the values of $\partial_y \epsilon$ in 0, r and L are gauge independent and this renders $\mathcal{L}_{eff}$ free from any gauge ambiguities.

3 Scalars Kinetic Energy

3.1 The compact case

In this section we will focus on the part of the effective Lagrangian involving the scalars and concentrate on the dilaton and radion kinetic term coefficients $\mathcal{K}_1$ and $\mathcal{K}_2$. In particular we are interested in the cases when the radion becomes a ghost field, *i.e.* $\mathcal{K}_2 < 0$. Firstly, it is trivial to obtain the dilaton kinetic term $\mathcal{K}_1$ by carrying out the first integral of (3.15):

$$\mathcal{K}_1 = 3c^2 M^3 \left[\frac{a^{-2}(r) - 1}{k_1} + \frac{a^{-2}(L) - a^{-2}(r)}{k_2}\right] \tag{3.18}$$

It turns out that for any possible values of k_1, k_2 and r, L the above quantity is positive definite. The radion kinetic term on the other hand is more involved. Integrating (3.12) we get the radion wavefunction for the two regions ($y > 0$):

$$B(y) = \begin{cases} c_1\, a^{-2} + 2k_1\, \partial_y \epsilon\, a^2 & , 0 < y < r \\ c_2\, a^{-2} + 2k_2\, \partial_y \epsilon\, a^2 & , r < y < L \end{cases} \tag{3.19}$$

where c_1 and c_2 are integration constants. The orbifold boundary conditions demand that $\partial_y \epsilon(0) = \partial_y \epsilon(L) = 0$ since ϵ is an even function of y. From the non-mixing conditions for radion and dilaton (3.13) and the continuity of B we are able to determine c_2 and $\partial_y \epsilon(r)$ as the following:

$$c_2 = c_1 \frac{k_2}{k_1} \, \frac{a^2(r) - 1}{\left(\frac{a(r)}{a(L)}\right)^2 - 1} \tag{3.20}$$

$$\epsilon'(r) = \frac{c_1 k_2}{2k_1(k_2 - k_1)a^4(r)} \left[\frac{k_1}{k_2} - \frac{a^2(r) - 1}{\left(\frac{a(r)}{a(L)}\right)^2 - 1} \right] \tag{3.21}$$

Therefore, the values of the radion wavefunction B at the brane positions are given by the following expressions:

$$B(0) = c_1 \tag{3.22}$$

$$B(r) = \frac{c_1 k_2}{(k_2 - k_1)} \frac{1 - a^2(L)}{a^2(L)\left[\left(\frac{a(r)}{a(L)}\right)^2 - 1\right]} \tag{3.23}$$

$$B(L) = \frac{c_1 k_2}{k_1} \frac{a^2(r) - 1}{a^2(L)\left[\left(\frac{a(r)}{a(L)}\right)^2 - 1\right]} \tag{3.24}$$

Thus, we can carry out the second integral on (3.15) to find the radion kinetic term coefficient:

$$\begin{aligned} \mathcal{K}_2 &= 3M^3 \left[\left(\frac{1}{k_1} - \frac{1}{k_2} \right) B^2(r)\, a^2(r) + \frac{1}{k_2}\, B^2(L)\, a^2(L) - \frac{1}{k_1}\, B^2(0) \right] \\ &= \frac{3M^3 c_1^2}{k_1} \left\{ \frac{k_2}{(k_2 - k_1)} \frac{a^2(r)(a^2(L) - 1)^2}{a^4(L)\left[\left(\frac{a(r)}{a(L)}\right)^2 - 1\right]^2} + \frac{k_2}{k_1} \frac{(a^2(r) - 1)^2}{a^2(L)\left[\left(\frac{a(r)}{a(L)}\right)^2 - 1\right]^2} - 1 \right\} \end{aligned} \tag{3.25}$$

The above quantity is not positive definite. In particular, it turns out that it is positive whenever the intermediate brane has positive tension and negative whenever the intermediate brane has negative tension. This result is graphically represented in Fig.3.2 where the (k_1, k_2) plane is divided in two regions.

3.2 The non-compact limit

It is instructive to discuss the decompactification limit in which the third brane is sent to infinity, *i.e.* $L \to +\infty$. To examine this limit we distinguish two cases:



Figure 3.2: Sign of the radion kinetic term in the (k_1, k_2) plane. In the regions A, B', C' the moving brane has positive tension and the radion positive kinetic energy. In the regions B, C, A' the moving brane has negative tension and the radion negative kinetic energy. We show the GRS line for the non-compact case. The dashed line corresponds to $k_1 = k_2$, *i.e.* a *tensionless* moving brane.

The case $\mathbf{k_2} > 0$

In this case we have $a(\infty) \propto \lim_{L\to\infty} e^{-k_2 L} = 0$. The dilaton kinetic term is trivial since from (3.18) we obtain $\mathcal{K}_1 \to \infty$. In other words the dilaton decouples from the four dimensional effective theory and the condition of absence of kinetic mixing between the scalars (3.13) plays no role. The radion kinetic term coefficient can be read off from (3.25):

$$\mathcal{K}_2 = \frac{3M^3 c_1^2}{k_1} \left[e^{2k_1 r} \frac{k_2}{(k_2 - k_1)} - 1 \right] \tag{3.26}$$

This result agrees with the computation of [32]. It is easy to see that the radion has positive kinetic energy when the moving brane has positive tension and negative kinetic energy when the tension is negative. Indeed, for $0 < k_1 < k_2$, or for $k_2 > 0$ and $k_1 < 0$ we have a positive brane and positive kinetic energy. On the other hand, for $0 < k_2 < k_1$ we have a negative tension brane and negative kinetic energy. In the limit $k_2 \to 0$ we get the GRS model with negative kinetic energy as in [32].

The case $\mathbf{k_2} < 0$

In this case $a(\infty) \propto \lim_{L\to\infty} e^{-k_2 L} \to \infty$. This time the dilaton plays in the game since

$\mathcal{K}_1$ is finite as seen from (3.18) and has the value:

$$\mathcal{K}_1 = \frac{3M^3\, c^2}{2k_1|k_2|}\left[e^{2k_1 r}(|k_2| + k_1) - |k_2|\right] \tag{3.27}$$

which is manifestly positive definite. The presence of dilaton mode is somewhat surprising since it describes the fluctuations of the overall size of the system which is infinite. Something similar happens in the Karch-Randall (KR) model [43] that has a dilaton mode although it is non-compact. The dilaton is a remnant of the decompactification process of the $'' + +''$ model [42] which we will examine in the following chapter and enters in the game because the inverse of the warp factor is normalizable.

The radion kinetic term coefficient can be obtained from taking the limit $L \to +\infty$ in (3.25). We get

$$\mathcal{K}_2 = \frac{3M^3\, c_1^2}{k_1}\left[e^{-2k_1 r}\,\frac{k_2}{(k_2 - k_1)} - 1\right] \tag{3.28}$$

The same considerations for the compact case applies here. When $k_1 < k_2 < 0$ we have a positive tension brane and a positive kinetic energy. On the other hand when $k_2 < k_1 < 0$, or for $k_2 < 0$ and $k_1 > 0$ we have a negative tension brane and a negative kinetic energy. In the GRS limit $k_2 \to 0$, the radion has negative kinetic energy.

4 Gravity on the branes

In this section we will study how the moduli and graviton(s) couple to matter confined on the branes. For simplicity we will study the non-compact case. For this purpose, we consider a matter Lagrangian $\mathcal{L}_m(\Phi_j, \hat{G}^{(i)})$, where we denote generically with Φ_j matter fields living on the branes; $\hat{G}^{(i)}$ is the induced metric on the i-th brane. Our starting point is the following action:

$$\begin{aligned}\mathcal{S} = \int d^4x dy\,\sqrt{-G}\,[2M^3 R - \Lambda(y)] - \sum_i \int d^4x\sqrt{-\hat{G}^{(i)}}V_i \\ + \sum_i \int d^4x dy\sqrt{-\hat{G}^{(i)}}\mathcal{L}_m(\Phi_j, \hat{G}^{(i)})\end{aligned} \tag{3.29}$$

We have already calculated the effective action for the gravity sector for the perturbation (3.10) in the previous chapter. It is useful to decompose the perturbation $\tilde{h}(x,z)$ in terms of a complete set of eigenfunctions[2] $\tilde{\Psi}^{(n)}(z)$ of the graviton kinetic operator (a suitable four dimensional gauge fixing like de Donder is understood):

---

[2] We denote by $\tilde{\Psi}(z)$ the gravitons wavefunctions in the conformal frame in contrast to the previous chapter graviton wavefunction $\Psi(z)$. Their relation is $\Psi(z) = a^2\tilde{\Psi}(z)$.

$$\tilde{h}(x,z)=\sum_n \tilde{\Psi}^{(n)}(z)\,h^{(n)}_{\mu\nu}(x)\,+\int dm\,\tilde{\Psi}(y,m)\,h^{(m)}_{\mu\nu}(x) \tag{3.30}$$

having taken into account both the discrete and the continuum part of the spectrum. The effective four dimensional Lagrangian has the following form:

$$\begin{aligned}\mathcal{L}=&\ \mathcal{L}_m(\Phi^c_j,\eta)+2M^3\sum_n^{*}\mathcal{L}_{PF}(h^{(n)}(x))+\mathcal{K}_1\,f_1\Box f_1+\mathcal{K}_2\,f_2\Box f_2\\ &-\sum_i\sum_n^{*}\frac{\tilde{\Psi}^{(n)}(y_i)}{2}h^{(n)}_{\mu\nu}(x)T^{\mu\nu}_{(i)}-\sum_i\frac{Q(y_i)}{2}f_1T_{(i)}-\sum_i\frac{B(y_i)}{2}f_2T_{(i)}\end{aligned} \tag{3.31}$$

The matter fields have been rescaled $\Phi_j\to\Phi^c_j$, to make them canonically normalized and the energy momentum tensor $T^{\mu\nu}_{(i)}$ for matter localized on the i-th brane is defined with respect to the rescaled fields Φ^c_j. By construction the induced background metric on the branes is the flat four dimensional Minkowski metric. The asterix denotes that the sum has to be converted into an integral for the continuum part of the spectrum. Finally, defining the canonically normalized fields:

$$\bar{h}^{(n)}_{\mu\nu}(x)=\sqrt{2M^3}h^{(n)}_{\mu\nu}(x)\,,\quad \bar{f}_1=\sqrt{2|\mathcal{K}_1|}f_1\,,\quad \bar{f}_2=\sqrt{2|\mathcal{K}_2|}f_2 \tag{3.32}$$

the Lagrangian reads:

$$\begin{aligned}\mathcal{L}=&\ \mathcal{L}_m(\Phi^c_j,\eta)+\sum_n^{*}\mathcal{L}_{PF}(\bar{h}^{(n)}(x))+\frac{1}{2}\,\bar{f}_1\Box\bar{f}_1+\frac{1}{2}\,\mathrm{sgn}(\mathcal{K}_2)\,\bar{f}_2\Box\bar{f}_2\\ &-\sum_i\sum_n^{*}\frac{\tilde{\Psi}^{(n)}(y_i)}{2\sqrt{2M^3}}\,\bar{h}^{(n)}_{\mu\nu}(x)\,T^{\mu\nu}_{(i)}-\sum_i\frac{Q(y_i)}{2\sqrt{2\mathcal{K}_1}}\,\bar{f}_1\,T_{(i)}-sum_i\frac{B(y_i)}{2\sqrt{2|\mathcal{K}_2|}}\,\bar{f}_2\,T_{(i)}\end{aligned} \tag{3.33}$$

Thus, the dilaton, the radion and the graviton (whenever it is normalizable) have the following couplings respectively:

$$\mathcal{C}_D=\frac{Q(y_{br})}{2\sqrt{2\mathcal{K}_1}}\,,\qquad \mathcal{C}_R=\frac{B(y_{br})}{2\sqrt{2|\mathcal{K}_2|}}\,,\qquad \mathcal{C}_G=\frac{\tilde{\Psi}^{(0)}(y_{br})}{2\sqrt{2M^3}} \tag{3.34}$$

For convenience we will define $\mathcal{C}_N\equiv\frac{\sqrt{|k_1|}}{2\sqrt{2M^3}}$ and write the above quantities as:

$$\mathcal{C}^{(i)}_D=\mathcal{C}_N\frac{g^{(i)}_D}{\sqrt{3}}\,,\qquad \mathcal{C}^{(i)}_R=\mathcal{C}_N\frac{g^{(i)}_R}{\sqrt{3}}\,,\qquad \mathcal{C}_G=\mathcal{C}_N\frac{\tilde{\Psi}^{(0)}(y_{br})}{\sqrt{|k_1|}} \tag{3.35}$$



Figure 3.3: The function $\sigma(y) = -\log[a(y)]$ for all possible combinations of k_1, k_2. The regions are named in accordance with the Fig.3.2 phase diagram.

where the dimensionless radion couplings are:

$$g_D^{(0)} = \frac{\theta(-k_2)}{\sqrt{\left|e^{2k_1 r \frac{k_1+|k_2|}{|k_2|}} - 1\right|}} \quad , \qquad g_D^{(r)} = e^{2k_1 r} g_D^{(0)} \tag{3.36}$$

$$g_R^{(0)} = \frac{1}{\sqrt{\left|e^{2k_1 r \mathrm{sgn}(k_2) \frac{k_2}{k_2-k_1}} - 1\right|}} \quad , \qquad g_R^{(r)} = e^{2k_1} \frac{k_2}{k_2 - k_1} g_R^{(0)} \tag{3.37}$$

We should note at this point that there are certain cases as we will see in the following that multigravity is realized and four dimensional gravity in intermediate distances is due to more than one mode. In that case we will denote by $\mathcal{C}_G$ the coupling of the "effective zero mode" even though there might not be a genuine zero mode at all.

We will now discuss how these couplings behave is the six distinct combinations of k_1 and k_2 shown in Fig.3.3.

Region A

In this case, we have the non-compact system of two positive tension branes discussed in [57,32,26]. The volume of the extra dimension is finite and therefore we have a normalizable graviton zero mode. The KK tower will be continuum and its coupling to matter on either branes will be exponentially suppressed for reasonably large r as can be easily seen by the analysis of [26]. The graviton wavefunction in the conformal gauge is constant, *i.e.* $\tilde{\Psi}^{(0)}(y) = A$. Thus the universal coupling of the zero mode will be $\mathcal{C}_G = \mathcal{C}_N$.

The dilaton in this region will simply be absent, or equivalently it will have zero coupling $\mathcal{C}_D = 0$ to the branes because $\mathcal{K}_1 \to \infty$. The radion in this region has positive kinetic term and from (3.37) we find that its coupling to matter on the $y = 0$ and $y = r$ branes is bounded as following (see Fig.3.4):

$$0 \leq g_R^{(0)} \lesssim e^{-k_1 r}\,, \qquad g_R^{(r)} \gtrsim e^{k_1 r} \tag{3.38}$$

The radion on the central brane is always weakly coupled and decouples in the tensionless moving brane limit. On the other hand the radion is always strongly coupled on the moving brane and diverges in the tensionless limit where $k_2 \to k_1$.

Region B

The volume of the extra dimension is still finite and therefore we have a normalizable graviton zero mode. When $k_1 = k_2$ the brane is tensionless and as soon as $k_2 < k_1$ the KK continuum starts to develop a “resonance”. This “resonance” is initially rather broad as $k_2 \to k_1$ and its width decreases until it coincides with the GRS width $\Gamma \sim k_1 \exp(-3k_1 r)$ as $k_2 \to 0$. In this region four dimensional gravity at intermediate distances is the net effect of the massless graviton and the lower part of the KK continuum that contributes more and more as $k_2 \to 0$.

The dilaton is again absent, $\mathcal{C}_D = 0$, since still we have $\mathcal{K}_1 \to \infty$. The radion in this region is a ghost field, to compensate for the presence of the extra polarization states of the contributing massive gravitons, and its coupling to the branes is bounded as (see Fig.3.4):

$$0 \leq g_R^{(0)} \leq 1\,, \qquad g_R^{(r)} \leq 0 \tag{3.39}$$

The radion on the central brane interpolates between the decoupling limit of the tensionless moving brane to the GRS limit where it couples with strength equal to the one of the graviton. On the other hand, the radion coupling on the moving brane interpolates between the infinitely strong limit of the tensionless brane (strictly speaking there is no brane in $y = r$ and we are left with RS2) to the GRS limit where it decouples.

Region C

In this region we have a infinite analogue of the $''+-+''$ model with no normalizable zero mode present. The graviton spectrum for $k_2 \to 0$ is approximately equally spaced and there is a resonance which coincides with the GRS width and gets more and more broad as k_2 gets more negative. Soon enough a special light state is singled out as in the $''+-+''$ model, a behaviour that persists for all values of k_2 in this region.

From the above behaviour we deduce that four dimensional gravity at intermediate distances is generated by the lower part of the discrete spectrum as $k_2 \to 0$, whereas only by the special state for all other values of k_2.

The dilaton in this region is present and its coupling to the branes is bounded as following (see Fig.3.4):

$$0 \leq g_D^{(0)} \lesssim e^{-k_1 r}\,, \qquad 0 \leq g_D^{(r)} \lesssim e^{k_1 r} \tag{3.40}$$

It is always weakly coupled to matter on the central brane and becomes strongly coupled but saturated on the moving brane.

The radion in this region is again a ghost field to cancel the unwanted extra massive graviton polarization states and its coupling to the branes is bounded as (see Fig.3.4):

$$g_R^{(0)} \approx 1\,, \qquad 0 \leq g_R^{(r)} \lesssim e^{k_1 r} \tag{3.41}$$

On the central brane it couples always with strength equal to the one of the "effective graviton" and on the moving brane it interpolates between the decoupling limit of the GRS case to a strongly coupled region with saturated coupling as the tension of the second brane gets infinite.

Region A'

In this region we have no normalizable zero mode and the two negative tension brane system resembles an inverted version of the RS2 model. The KK spectrum is discrete and all excitations lie above the characteristic curvature scale k_1. Thus the low energy effective theory does not have four dimensional gravity at all.

The dilaton in this system is present and the coupling to the branes is approximately constant (see Fig.3.4):

$$g_D^{(0)} \approx 1\,, \qquad g_D^{(r)} \approx 1 \tag{3.42}$$

The radion field is a ghost and its coupling to matter on the branes is bounded as (see Fig.3.4):

$$g_R^{(0)} \gtrsim e^{-3|k_1|r}\,, \qquad 0 \leq g_R^{(r)} \lesssim e^{-2|k_1|r} \tag{3.43}$$

The radion on the central negative tension brane is always weakly coupled and vanishes in the limit of $k_2 \to k_1 = -|k_1|$ of the tensionless moving brane. On the other hand the coupling on the moving brane is bounded from below in the limit of infinite negative tension brane and diverges at the limit of the tensionless brane.



Figure 3.4: The dimensionless couplings of the radion (upper) and the dilaton (lower) to matter on the $y = 0$ or the $y = r$ branes. The left diagrams correspond to $k_1 > 0$ while the right ones for $k_1 < 0$. The regions $A,\ B, \ldots$ are in accordance with the ones of the phase diagram in Fig.3.2. The diagrams are not in scale.

Region B'

In this region there is again no normalizable zero mode and the system of the negative and positive tension branes still resembles the inverted RS2 model. The KK spectrum is almost identical with the one of the previous region except for the limit $k_2 \to 0$ when the spectrum drops below the curvature scale k_1 and a continuum develops.

The dilaton coupling to the branes decreases from the constant value of the previous region, to zero as $k_2 \to 0$ (see Fig.3.4):

$$0 \le g_D^{(0)} \lesssim 1\,, \qquad 0 \le g_D^{(r)} \lesssim e^{-2|k_1|r} \tag{3.44}$$

The radion has positive kinetic energy and its coupling to the branes is bounded as following (see Fig.3.4):

$$0 \leq g_R^{(0)} \leq 1\,, \qquad g_R^{(r)} \leq 0 \tag{3.45}$$

It is weakly coupled in the central brane and the coupling interpolates between zero from the previous region to one in the inverted GRS case where $k_2 \to 0$. On the second brane it is divergent as $k_2 \to k_1 = -|k_1|$ and vanishes as $k_2 \to 0$.

Region C'

In this region we again have a system of a negative and a positive tension brane, but gravity can be localized on the moving positive tension brane. There is a normalizable zero mode that mediates four dimensional gravity and a KK continuum with suppressed couplings on the branes.

The dilaton in this region will be absent, or equivalently it will have zero coupling $\mathcal{C}_D = 0$ to the branes because $\mathcal{K}_1 \to \infty$.

The radion will have again positive kinetic term and its coupling will be bounded as following (see Fig.3.4):

$$0 \leq g_R^{(0)} \lesssim e^{-2|k_1|r}\,, \qquad g_R^{(r)} \approx 1 \tag{3.46}$$

It is approximately constant on the central brane with strength equal to the one of the graviton and is weakly coupled to the moving brane with coupling interpolating between zero in the inverted GRS case with $k_2 \to 0$ to a certain value as the tension of the moving brane becomes infinitely large.

5 Review of sum rules

In this section we will review the sum rules presented in [41] in order to discuss the dilaton stabilization in the presence of bulk dynamics of arbitrary minimally coupled bulk scalar fields. We will concentrate as in [41] in the case where the background metric of the five dimensional spacetime can be written in the form:

$$ds^2 = a(y)^2 g_{\mu\nu}(x) dx^\mu dx^\nu + dy^2 \tag{3.47}$$

with $g_{\mu\nu}(x)$ a general background four dimensional metric and $a(y)$ a generic warp factor. We should stress here that this is not the most general choice of metric in five dimensions as we have explicitly assumed that the all four dimensional sections have the same conformal geometry. The particular choice $g_{\mu\nu} = \eta_{\mu\nu}$ is valid for the models which we have discussed so far.

We can now consider an arbitrary number of minimally coupled scalar bulk fields $\Phi^I(x,y)$ with internal metric $\mathcal{G}_{IJ}$ and arbitrary bulk potential $V(\Phi)$ (which includes bulk cosmological constant), coupled to an again arbitrary number of branes with brane potential $\lambda_i(\Phi)$ (which again includes the brane tensions). The action describing the above system is the following:

$$\mathcal{S} = \int d^4x dy \sqrt{-G}\left(2M^3 R - \frac{1}{2}\mathcal{G}_{IJ}\partial_M\Phi^I\partial^M\Phi^J - V(\Phi) - \sum_i \lambda_i(\Phi)\delta(y-y_i)\frac{\sqrt{-\hat{G}^{(i)}}}{\sqrt{-G}}\right) \tag{3.48}$$

where $\hat{G}^{(i)}_{\mu\nu}$ is the induced metric on the brane and M the fundamental five dimensional scale. The Einstein equations arising from the above metric can be written in the form:

$$4M^3 R^\mu_\mu = -\frac{1}{3}T^\mu_\mu - \frac{4}{3}T^5_5 \tag{3.49}$$

$$4M^3 R^5_5 = -\frac{1}{3}T^\mu_\mu + \frac{2}{3}T^5_5 \tag{3.50}$$

where the energy-momentum tensor components are:

$$T^\mu_\mu = -\partial_\mu\Phi\cdot\partial^\mu\Phi - 2\Phi'\cdot\Phi' - 4V(\Phi) - 4\sum_i \lambda_i(\Phi)\delta(y-y_i) \tag{3.51}$$

$$T^5_5 = -\frac{1}{2}\partial_\mu\Phi\cdot\partial^\mu\Phi + \frac{1}{2}\Phi'\cdot\Phi' - V(\Phi) \tag{3.52}$$

with the indices in the above formulas raised and lowered by $G_{\mu\nu} = a(y)^2 g_{\mu\nu}(x)$ and where dot product denotes contraction with the internal metric $\mathcal{G}_{IJ}$. Since we are interested in a background configuration, the $\partial_\mu\Phi\cdot\partial^\mu\Phi$ terms can be dropped. The Ricci tensor is easily calculated to be:

$$R^\mu_\mu = a^{-2}R_g - 12(a')^2a^{-2} - 4a''a^{-1} \tag{3.53}$$

$$R^5_5 = -4a''a^{-1} \tag{3.54}$$

If we now consider the function $(a^n)''$ with n an arbitrary real number, its integral around the compact extra dimension is zero. Using (3.49), (3.50), (3.53), (3.54) we arrive at an infinite number of constraints [41]:

$$\oint dy\ a^n(T^\mu_\mu + (2n-4)T^5_5) = 4M^3(1-n)R_g \oint dy\ a^{n-2} \tag{3.55}$$

As it is obvious, these constraints are a natural consequence of the equations of motion and are totally equivalent to them. It is straightforward to see for example that they are satisfied in the RS1 model [16] and the bigravity/multigravity models [21, 26, 42]. We will single out three constraints which we will be important for the subsequent discussion, namely the ones for $n = 0, 1, 2$:

$$\oint dy\ (T^\mu_\mu - 4T^5_5) \ = \ 4M^3 R_g \oint dy\ a^{-2} \tag{3.56}$$

$$\oint dy\ a(T^\mu_\mu - 2T^5_5) \ = \ 0 \tag{3.57}$$

$$\oint dy\ a^2 T^\mu_\mu \ = \ -4M^3 R_g \oint dy \tag{3.58}$$

One could also use the above constraints for non-compact models, but should be careful that the above derivation makes sense. For the RS2 model [15], all constraints are valid for $n \geq 0$. For the GRS model [22] only the $n = 0$ constraint is valid and for the KR model [43] (see next chapter) all constraints are valid for $n \leq 0$.

6 Dilaton in warped backgrounds

We now consider the perturbation related to the overall size of the compact system, namely the dilaton. As previously noted in the chapter, we do not consider the KK-tower of the dilaton since these states can be gauged away by five dimensional diffeomorphisms. The general form of the metric for the physical radion perturbations that do not mix with the graviton(s) is given in [32, 33]. For the dilaton, the ansatz is rather simple and following [34] we write it in the form[3]:

$$ds^2 = e^{-a(y)^{-2}\gamma(x)} a(y)^2 g_{\mu\nu}(x) dx^\mu dx^\nu + \left(1 + a(y)^{-2}\gamma(x)\right)^2 dy^2 \tag{3.59}$$

Substituting the above metric in the action (3.48) (see Appendix B for analytic formulas), integrating out total derivatives, throwing out γ-independent parts and keeping terms up to quadratic order, we get:

$$\mathcal{S} = \int d^4x dy \sqrt{-g} \left\{ -\frac{1}{2}(6M^3 a^{-2}) g^{\mu\nu} \gamma_{,\mu} \gamma_{,\nu} + \mathcal{L}_1 \gamma - \frac{1}{2}\mathcal{L}_2 \gamma^2 \right\} \tag{3.60}$$

---

[3] We are interested only in the quadratic in $\gamma(x)$ Lagrangian derived from the above metric, so only the part of the metric up to linear order in $\gamma(x)$ is important. However, we prefer to keep the same notation as [34] and do the calculation with (3.59). The result, of course, for the quadratic Lagrangian would be the same if we used the metric up to linear order in $\gamma(x)$.

with

$$\mathcal{L}_1 = 2M^3(4(a')^2 + 16a''a) + \frac{3}{2}a^2\Phi' \cdot \Phi' + a^2V(\Phi) + 2a^2\sum_i \lambda_i(\Phi)\delta(y-y_i) \quad (3.61)$$

$$\mathcal{L}_2 = 2M^3(a^{-2}R_g - 32(a')^2a^{-2} + 32a''a^{-1}) + 5\Phi' \cdot \Phi' + 4\sum_i \lambda_i(\Phi)\delta(y-y_i) \quad (3.62)$$

At this point, let us work out the integral over the extra dimension of the tadpole term $\mathcal{L}_1$ of the Lagrangian. This gives:

$$\oint dy\mathcal{L}_1 = -\frac{1}{2}\oint dy\ a^2\left[T^\mu_\mu - 2T^5_5 - 16M^3\left(\frac{(a')^2}{a^2} + 4\frac{a''}{a}\right)\right] \quad (3.63)$$

where we used the energy-momentum tensor components found in the previous section with respect to the unperturbed background metric (3.51), (3.52). We can further simplify this quantity if we use the equations (3.49), (3.50), (3.53), (3.54) which hold for the background metric. The resulting expression is:

$$\oint dy\mathcal{L}_1 = \frac{1}{6}\oint dy\ a^2\left[T^\mu_\mu + 4M^3a^{-2}R_g\right] \quad (3.64)$$

which is exactly zero because of the $n = 2$ constraint (3.58). This result should have been expected since the perturbation (3.59) is bound to extremize the effective potential when one evaluates the action using the background equations of motion. It is worth mentioning here that in the case that the warp factor is effectively constant ($a \approx 1$), as it was assumed in [58], the condition that the expression (3.63) vanishes, is identical with the $n = 1$ constraint (3.57).

Our next task is to read off the mass of the dilaton from the action functional. For this reason we define the canonically normalized dilaton field with mass dimension one $\bar{\gamma}^2 = \left(6M^3\oint dya^{-2}\right)\gamma^2 \equiv A\gamma^2$. Then the mass of the canonical dilaton $\bar{\gamma}$ is:

$$m^2 = \frac{1}{A}\oint dy\mathcal{L}_2 \quad (3.65)$$

After a lot of simplifications using the relations (3.49)-(3.54) we obtain:

$$m^2 = -\frac{1}{3A}\oint dy(10M^3a^{-2}R_g + \Phi' \cdot \Phi' + 4\sum_i \lambda_i(\Phi)\delta(y-y_i)) \quad (3.66)$$

We can further simplify the expression using the $n = 0$ constraint (3.56) and get a more suggestive result:

$$m^2 = \frac{1}{A} \oint dy(\Phi' \cdot \Phi' - 2M^3 a^{-2} R_g) \tag{3.67}$$

or equivalently,

$$m^2 = -\frac{1}{A} \left\{ \sum_i \lambda_i(\Phi) + 3M^3 R_g \oint dy a^{-2} \right\} \tag{3.68}$$

From the second expression it is clear that we cannot have a massive dilaton if the sum of the effective tensions of the branes $\lambda_i(\Phi)$ is exactly zero and at the same time they are kept flat. This is the same conclusion that appeared in [41] regarding the Goldberger-Wise mechanism [59] in the RS1 scenario. Regarding the latter mechanism this is not a surprise, since promoting the Goldberger-Wise to an exact solution means that either the scalar field energy will generate a four dimensional cosmological constant, or if one insists on having flat branes, this will introduce a new and different fine tuning of the tensions which gives a mismatch of brane tensions. Moreover, the absence of a tachyonic mass will guarantee the stabilization of the overall size of any system with the above characteristics. By (3.67), (3.68) we get two equivalent conditions:

$$\oint dy(\Phi' \cdot \Phi' - 2M^3 a^{-2} R_g) \quad > \quad 0 \tag{3.69}$$

$$\sum_i \lambda_i(\Phi) + 3M^3 R_g \oint dy a^{-2} \quad < \quad 0 \tag{3.70}$$

If one wishes to have flat branes, then the sum of brane tensions should be negative or equivalently one should have a non constant (in y) scalar field configuration. In the case that the above expressions (and thus the mass) vanish, one should look for the higher orders of the effective potential to examine the stability of the system.

Finally, we should note that the formulas (3.67), (3.68) are valid even for non-compact models whenever the dilaton mode is normalizable. This happens for example in the KR model [43] and one can find from the above expressions the mass of the dilaton, in agreement with [60]. In that case, the dilaton mode cannot be attributed to the fluctuation of the overall size of the system, but can be understood to be a remnant mode if we start with the compact $'' + +''$ system [42] and send one of the branes to infinity (*i.e.* after decompactifying the system). We will see this possibility in the next chapter.

Chapter 4

AdS_4 brane multigravity models in five dimensions

As we have seen clearly in the previous two chapters, the flat brane multigravity models in five dimensions necessarily have moving negative tension branes. These in turn have corresponding radions with negative kinetic energy which are disastrous for the theory. It is clearly desirable to have a model with only positive tension branes. In this chapter we will demonstrate how one can obtain such a model in five dimensions. The price that one has to pay is that the branes are no longer flat, but instead AdS_4. A two brane model with only positive tension branes was already known in [63,64,65,61,58,62,66] and exhibited a characteristic bounce of the warp factor. We know from chapter 2 that a model that has a bounce in the warp factor is bound to have bigravity, something that we will explicitly show for this particular model in the present chapter. Furthermore, we can construct a crystal model with an infinite array of positive tension branes, but we cannot reproduce the quasi-localized model because the presence of the cosmological constant on the branes prevents the warp factor from being asymptotically constant.

Since there is no ghost modulus in these configurations one may worry about the structure of the graviton propagator at intermediate distances. As we will show explicitly, the AdS_4 brane configurations can evade this problem because of an unusual property of the graviton propagator in AdS background. We will show in the second part of this chapter that the van Dam-Veltman-Zakharov discontinuity can in certain cases be circumvented and a particular example is the the $'' + +''$ model. In more detail, the troublesome additional polarizations are very weakly coupled and do not disturb the normal structure of the graviton propagator.

Nevertheless, the construction that we obtain faces phenomenological difficulties because of the remnant negative cosmological constant on the brane. This sets an horizon scale and unfortunately the Compton wavelength of the light state of the system lies exponentially far from this horizon. This does not change even if we consider



Figure 4.1: The $''++''$ bigravity model with two AdS_4 $''+''$ branes at the fixed points of a S^1/Z_2.

a highly asymmetric version of this model. As a result, this bigravity model makes no predictions for observable deviations from Newtonian gravity at ultra-large distances. Furthermore, the very fact that the remnant cosmological constant is negative is observationally disfavoured.

1 The $''++''$ bigravity model

The model consists of two three-branes with tensions V_1 and V_2 respectively, in an AdS_5 space with five dimensional cosmological constant $\Lambda < 0$. The fifth dimension has the geometry of an orbifold and the branes are located at its fixed points, *i.e.* $L_0 = 0$ and $L_1 = L$ (see Fig.4.1). Due to orbifolding, we can restrict ourselves to the region $0 \leq y \leq L$. Firstly, we find a suitable vacuum solution. The action of this setup is:

$$\mathcal{S}_G = \int d^4x \int_{-L}^{L} dy \sqrt{-G}[2M^3R - \Lambda] - \sum_i \int_{y=L_i} d^4x V_i \sqrt{-\hat{G}^{(i)}} \tag{4.1}$$

where $\hat{G}^{(i)}_{\mu\nu}$ is the induced metric on the branes. The Einstein equations that arise from this action are:

$$R_{MN} - \frac{1}{2}G_{MN}R = -\frac{1}{4M^3}\left(\Lambda G_{MN} + \sum_i V_i \frac{\sqrt{-\hat{G}^{(i)}}}{\sqrt{-G}} \hat{G}^{(i)}_{\mu\nu}\delta^{\mu}_{M}\delta^{\nu}_{N}\delta(y - L_i)\right) \tag{4.2}$$

In order to find a specific form of the equations of motion we need to write a metric ansatz which will take in account the spacetime symmetries of the three-brane. Since we would like not to restrict our model to flat solutions on the branes, we should make a choice which will let us interpolate between the maximally symmetric space-times in four dimensions, *i.e.* de-Sitter, Minkowski and Anti-de-Sitter. The metric ansatz that accomplishes this is the following:

$$ds^2 = \frac{a^2(y)}{(1 \mp \frac{H^2 x^2}{4})^2}\eta_{\mu\nu}dx^\mu dx^\nu + dy^2 \tag{4.3}$$

where $x^2 = \eta_{\mu\nu}x^\mu x^\nu$. It can be shown that the Ricci scalar for this metric is $R = \mp 12H^2$. From now on, the upper sign corresponds to AdS_4 space while the lower to dS_4 space and always $H^2 > 0$. Thus, this metric represents all maximally symmetric spaces. A straightforward calculation of the Einstein's equations gives us the following differential equations for $a(y)$:

$$\begin{aligned}(a'(y))^2 &= \pm H^2 - \frac{\Lambda}{24M^3}a^2(y) &(4.4)\\ a''(y) &= -\sum_i \frac{V_i}{12M^3}a(y)\delta(y-L_i) - \frac{\Lambda}{24M^3}a(y) &(4.5)\end{aligned}$$

By solving the above equations we find that the solution can be written in the form:

$$a(y) = \cosh(k|y|) + \frac{V_1 k}{\Lambda}\sinh(k|y|) \tag{4.6}$$

with

$$H^2 = \begin{cases} \frac{k^2}{\Lambda^2}(V_1^2k^2 - \Lambda^2) & , \frac{|\Lambda|}{k} < V_1 \ \text{ for } dS_4 \text{ branes} \\ 0 & , \frac{|\Lambda|}{k} = V_1 \ \text{ for flat branes} \\ \frac{k^2}{\Lambda^2}(\Lambda^2 - V_1^2k^2) & , \frac{|\Lambda|}{k} > V_1 \ \text{ for } AdS_4 \text{ branes} \end{cases} \tag{4.7}$$

where we have normalized $a(0) = 1$ and assumed $V_1 > 0$. Also we have defined $k \equiv \sqrt{\frac{-\Lambda}{24M^3}}$.

Additionally, in order to have this solution, the brane tensions V_1, V_2, the bulk cosmological constant $|\Lambda|$ and the position of the second brane L must be related through the equation:

$$\tanh(kL) = k|\Lambda|\frac{V_1 + V_2}{|\Lambda|^2 + k^2 V_1 V_2} \tag{4.8}$$

Let us now restrict ourselves to the case of AdS_4 spacetime on the two branes which will turn out to be the most interesting. In this case the condition $\frac{|\Lambda|}{k} > V_1$ must hold.

Hence, we can define $\tanh(ky_0) \equiv \frac{kV_1}{|\Lambda|}$ and write the solution in the form:

$$a(y) = \frac{\cosh(k(y_0 - |y|))}{\cosh(ky_0)} \tag{4.9}$$

from which it is clear that the warp factor has a minimum at $y = y_0$. From this point we can see the role of the AdS_4 on the branes, *i.e.* the role of the condition $\frac{|\Lambda|}{k} > V_1$. This condition allows us to have the bounce form of the warp factor and the second brane to have positive tension and give us, as we will see shortly, a phenomenology quite similar to the $'' + - +''$ bigravity model [21]. This can be easily seen from the (4.8) which relates the brane tensions and the distance between the branes. From this equation we indeed see that by placing the second brane after the minimum of the warp factor we can make the tension of the second brane positive and thus both branes that appear in the model have positive tension avoiding the problems associated with negative tension branes. In fact it is clear that the present model mimics the characteristics of the $'' + - +''$ bigravity model since what we effectively do is to reproduce the effect of the presence of a negative tension brane, *i.e.* the bounce form of the warp factor, with another mechanism allowing a negative four dimensional cosmological constant on the brane. Note that in the limit that $V_1 \rightarrow \frac{|\Lambda|}{k}$ (flat limit) the minimum of the warp factor tends to infinity and if we wish to have a second brane at some finite distance, it will necessarily have negative tension.

The relationship between the four dimensional effective fundamental scale M_* and the five dimensional fundamental scale M can be easily found by dimensional reduction to be:

$$M_*^2 = \frac{M^3}{k\cosh^2(ky_0)}\left[kL + \sinh(kL)\cosh(k(L - 2y_0))\right] \tag{4.10}$$

The above formula tells us that for finite L the compactification volume is finite and thus the zero mode is normalizable. In the case where we send the second brane to infinity, the compactification volume becomes infinite which indicates that the zero mode becomes non-normalizable. Note that M_* is not necessarily equal to the four dimensional Planck scale $M_{\rm Pl}$ since as will see shortly, at least for a sector of the parameter space of our model, gravity is the result not only of the massless graviton but also of an ultralight KK state.

The warp factor renormalizes the physical scales of the theory as in [16]. Thus, all mass parameters m_0 on a brane placed at the point y_{br} are rescaled as:

$$m = a(y_{br})m_0 \tag{4.11}$$

Hence, one can choose a distance between the two branes such that this rescaling leads to the creation of a desired mass hierarchy. Let us note at this point that this interbrane distance, and thus the resulting hierarchy, is dictated by the relative strength of the tensions of the two branes. This is not the case as the flat brane models where the hierarchy is to be fixed by some stabilization mechanism. The *AdS* brane system is self-stabilized and this can be easily seen by the fact that the dilaton in this case is massive. Indeed from (3.68) we obtain $m^2 = 4H^2 > 0$ which means that the system is truly stabilized.

However, since we consider non-flat solutions on the branes, we have to make sure that the four dimensional effective cosmological constant does not contradict present experimental and observational bounds. Recent experimental data favour a positive cosmological constant, nevertheless since zero cosmological constant is not completely ruled out it can be argued that also a tiny negative cosmological constant can be acceptable within the experimental uncertainties. The effective cosmological constant on the two branes is:

$$\Lambda_{4d} = -12H^2 M_*^2 = -\frac{12}{\cosh^2(ky_0)} k^2 M_*^2 \tag{4.12}$$

From the above formula we can see that we can make the cosmological constant small enough $|\Lambda_{4d}| \lesssim 10^{-120} M_{\rm Pl}^4$ if we choose large enough ky_0, *i.e.* $ky_0 \gtrsim 135$. This however will make observable deviations from Newtonian gravity at ultra-large scales impossible as we will see later.

To determine the phenomenology of the model we need to know the KK spectrum that follows from the dimensional reduction. This is determined by considering the linear fluctuations of the metric around the vacuum solution that we found above. We can write the metric perturbation in the form:

$$ds^2 = \left[a(y)^2 g_{\mu\nu}^{AdS} + h_{\mu\nu}(x,y)\right] dx^\mu dx^\nu + dy^2 \tag{4.13}$$

where $g_{\mu\nu}^{AdS}$ is the four dimensional AdS_4 metric. Here we have ignored the dilaton mode that has been calculated in the previous chapter. We expand the field $h_{\mu\nu}(x,y)$ into graviton and KK plane waves:

$$h_{\mu\nu}(x,y) = \sum_{n=0}^{\infty} h_{\mu\nu}^{(n)}(x)\Psi^{(n)}(y) \tag{4.14}$$

where we demand $\left(\nabla_\kappa \nabla^\kappa + 2H^2 - m_n^2\right) h_{\mu\nu}^{(n)} = 0$ and additionally $\nabla^\alpha h_{\alpha\beta}^{(n)} = h^{(n)\alpha}_{\ \ \alpha} = 0$.

The function $\Psi^{(n)}(y)$ will obey the second order differential equation:

$$\left\{-\frac{1}{2}\partial_y^2+\left(\frac{a'}{a}\right)^2+\frac{a''}{a}\right\}\Psi^{(n)}(y)=\frac{m_n^2}{2}\frac{1}{a^2}\Psi^{(n)}(y) \tag{4.15}$$

After a change of variables and a redefinition of the wavefunction the above equation reduces to an ordinary Schrödinger-type equation:

$$\left\{-\frac{1}{2}\partial_z^2+V(z)\right\}\hat{\Psi}^{(n)}(z)=\frac{m_n^2}{2}\hat{\Psi}^{(n)}(z) \tag{4.16}$$

where the potential is given by:

$$\begin{aligned} V(z)= & -\frac{9\tilde{k}^2}{8}+\frac{15\tilde{k}^2}{8}\frac{1}{\cos^2\left(\tilde{k}(|z|-z_0)\right)} \\ & -\frac{3k}{2}\left[\tanh(ky_0)\delta(z)+\frac{\sinh(k(L-y_0))}{\cosh(ky_0)}\delta(z-z_1)\right] \end{aligned} \tag{4.17}$$

with $\tilde{k}$ defined as $\tilde{k}\equiv\frac{k}{\cosh(ky_0)}$. The new variables and the redefinition of the wavefunction are related with the old ones by:

$$z\equiv\mathrm{sgn}(y)\frac{2}{\tilde{k}}\left[\arctan\left(\tanh(\frac{k(|y|-y_0)}{2})\right)+\arctan\left(\tanh(\frac{ky_0}{2})\right)\right] \tag{4.18}$$

$$\hat{\Psi}^{(n)}(z)\equiv\frac{1}{\sqrt{a(y)}}\Psi^{(n)}(y) \tag{4.19}$$

Thus, in terms of the new coordinates, the branes are at $z_{L_0}=0$ and z_L, with the minimum of the potential at $z_0=\frac{2}{\tilde{k}}\arctan\left(\tanh(\frac{ky_0}{2})\right)$. Also note that with this transformation the point $y=\infty$ is mapped to the finite point $z_\infty=\frac{2}{\tilde{k}}\left[\frac{\pi}{4}+\arctan\left(\tanh(\frac{ky_0}{2})\right)\right]$.

From now on we restrict ourselves to the symmetric configuration of the two branes with respect to the minimum z_0 (*i.e.* the first brane at 0 and the second at $2z_0$), since the important characteristics of the model appear independently of the details of the configuration. Thus, the model has been reduced to a quantum mechanical problem with δ-function potentials wells of the same weight and an extra smoothing term in-between (due to the AdS geometry). This gives the potential a double “volcano” form (see Fig.4.2). An interesting characteristic of this Schrödinger equation is that, as in the flat brane case, it has a “supersymmetric” form (see [67]) which guarantees that



Figure 4.2: The potential $V(z)$ in the $''++''$ bigravity model for the symmetric configuration. The δ-function wells correspond to $''+''$ branes.

it has no tachyonic modes.

Since the system is compact, *i.e.* for $z_L < z_\infty$, it gives rise to a normalizable massless zero mode, something that is expected since the volume of the extra dimension is finite. The zero mode wavefunction is given by:

$$\hat{\Psi}^{(0)}(z) = \frac{A}{[\cos(\tilde{k}(z_0 - |z|))]^{3/2}} \tag{4.20}$$

where the normalization factor A is determined by the requirement $\displaystyle\int_{-z_L}^{z_L} dz \left[\hat{\Psi}^{(0)}(z)\right]^2 = 1$, chosen so that we get the standard form of the Fierz-Pauli Lagrangian.

The form of the zero mode resembles the one of the zero mode of the $''+-+''$ model, *i.e.* it has a bounce form with the turning at z_0 (see Fig.4.3). In the case of the $''+-+''$ model the cause for this was the presence of the $''-''$ brane. In the present model considering AdS_4 spacetime on the branes we get the same effect.

In the case that we send the second brane to infinity (*i.e.* $z \to z_\infty$) the zero mode fails to be normalizable due to singularity of the wavefunction exactly at that point. This can be also seen from (4.10) which implies that at this limit M_* becomes infinite (*i.e.* the coupling of the zero mode becomes zero). Thus in this limit the model has no zero mode and all gravitational interactions must be produced by the ultralight first KK mode. The spectrum in this case was discussed by Randall and Karch in [43].

Considering the Schrödinger equation for $m \neq 0$ we can determine the wavefunctions of the KK tower. It turns out that the differential equation can be brought to a hypergeometric form, and hence the general solution is given in terms two hypergeometric functions:

$$\begin{aligned}\hat{\Psi}^{(n)} = \cos^{5/2}(\tilde{k}(|z|-z_0)) \Big[& C_1\ F(\tilde{a}_n, \tilde{b}_n, \tfrac{1}{2}; \sin^2(\tilde{k}(|z|-z_0))) \\ & + C_2\ |\sin(\tilde{k}(|z|-z_0))|\ F(\tilde{a}_n + \tfrac{1}{2}, \tilde{b}_n + \tfrac{1}{2}, \tfrac{3}{2}; \sin^2(\tilde{k}(|z|-z_0)))\Big]\end{aligned} \tag{4.21}$$

where

$$\begin{aligned}\tilde{a}_n &= \frac{5}{4} + \frac{1}{2}\sqrt{\left(\frac{m_n}{\tilde{k}}\right)^2 + \frac{9}{4}} \\ \tilde{b}_n &= \frac{5}{4} - \frac{1}{2}\sqrt{\left(\frac{m_n}{\tilde{k}}\right)^2 + \frac{9}{4}}\end{aligned} \tag{4.22}$$

with the same normalization convention as the zero mode. The boundary conditions (*i.e.* the jump of the wave function at the points $z = 0,\ z_L$) result in a 2×2 homogeneous linear system which, in order to have a non-trivial solution, should have vanishing determinant. In the symmetric configuration which we consider, this procedure can be simplified by considering even and odd functions with respect to the minimum of the potential z_0.

In more detail, the odd eigenfunctions obeying the boundary condition $\hat{\Psi}^{(n)}(z_0) = 0$ will have $C_1 = 0$ and thus the form:

$$\hat{\Psi}^{(n)} = C_2 \cos^{5/2}(\tilde{k}(|z|-z_0))|\sin(\tilde{k}(|z|-z_0))|\ F(\tilde{a}_n + \frac{1}{2}, \tilde{b}_n + \frac{1}{2}, \frac{3}{2}; \sin^2(\tilde{k}(|z|-z_0))) \tag{4.23}$$

On the other hand, the even eigenfunctions obeying the boundary condition $\hat{\Psi}^{(n)\ \prime}(z_0) = 0$ will have $C_2 = 0$ and thus the form:

$$\hat{\Psi}^{(n)} = C_1 \cos^{5/2}(\tilde{k}(|z|-z_0)) F(\tilde{a}_n, \tilde{b}_n, \frac{1}{2}; \sin^2(\tilde{k}(|z|-z_0))) \tag{4.24}$$

The remaining boundary condition is given by:

$$\hat{\Psi}^{(n)\ \prime}(0) + \frac{3k}{2}\tanh(ky_0)\hat{\Psi}^{(n)}(0) = 0 \tag{4.25}$$



Figure 4.3: The graviton (blue line), first (red line) and second (green line) KK states wavefunctions in the symmetric $''++''$ model. All wavefunctions are continuous at the position of the bounce. On the right the pattern of the spectrum is sketched.

and determines the mass spectrum of the KK states. From this quantization condition we get that the KK spectrum has a special first mode similar to the one of the $''+-+''$ bigravity model. For $ky_0 \gtrsim 5$ the mass of the first mode is given by the approximate relation:

$$m_1 = 4\sqrt{3}\ k\ e^{-2ky_0} \tag{4.26}$$

In contrast, the masses of the next levels, if we put together the results for even and odd wavefunctions, are given by the formula:

$$m_{n+1} = 2\sqrt{n(n+3)}\ k\ e^{-ky_0} \tag{4.27}$$

with $n = 1, 2, ...$

We note that the first KK state has a different scaling law with respect to the position of the minimum of the warp factor compared to the rest of the KK tower, since it scales as e^{-2ky_0} while the rest of the tower scales as e^{-ky_0}. Thus, the first KK state is generally much lighter than the rest of the tower. It is clear that this mass spectrum resembles the one of the $''+-+''$ bigravity model. The deeper reason for this is again the common form of the warp factor. In both cases the warp factor has a minimum due to its bounce form. The graviton wave function follows the form of the warp factor, *i.e.* it is symmetric with respect to z_0, while the wavefunction of the first KK state is antisymmetric in respect to z_0 (see Fig.4.3). The absolute values of the two wavefunctions are almost identical in all regions except near z_0 (the symmetric is nonzero while the antisymmetric is zero at z_0). The graviton wavefunction is suppressed by the factor $\cosh^{-2}(ky_0)$ at z_0 which brings its value close to zero for reasonable values of ky_0. Thus, the mass difference which is determined by the wavefunction near z_0 is expected to be generally very small, a fact which appears as the extra suppression factor e^{-ky_0} in the formula of m_1 in comparison with the rest of the KK tower.

In the case that we consider an asymmetric brane configuration, for example $z_L > 2z_0$, the spectrum is effectively independent of the position of the second brane z_L (considering $ky_0 \gtrsim 5$). Thus, even in the case that we place the second brane at z_∞, *i.e.* the point which corresponds to infinity in the y-coordinates, the spectrum is given essentially by the same formulas. This decompactification limit is exactly what has been studied in [43].

The coupling also of the first KK state to matter on the branes is very different from the one of the remaining tower. We can easily calculate it by the same procedure presented in chapter 2 considering the a minimally coupled to gravity matter Lagrangian. In the case of scale equivalent positive tension branes, the gravitons will couple to matter on them ($z = z_{br} = 0$ or L) with the following couplings:

$$a_n = \frac{\hat{\Psi}^{(n)}\left(z_{br}\right)}{2\sqrt{2M^3}} \tag{4.28}$$

In more detail we find that the zero mode and the first ultralight KK state couple with the same strength on the positive tension branes:

$$a_0 = a_1 = \frac{1}{2\sqrt{2}M_*} \tag{4.29}$$

while the coupling of the remaining tower is exponentially suppressed (it scales as e^{-ky_0}). The reason for the strong coupling of the first KK state in comparison with the remaining tower is readily understood because, being dominantly a bound state of the volcano potential on the positive tension brane, it is largely localized on it.

Exploiting the different mass scaling of the first KK relative to the rest we can ask whether it is possible to realize a bigravity scenario similar to that in $'' + - +''$ bigravity model. In that model by appropriately choosing the position of the negative tension brane, it was possible to make the first KK state have mass such that the corresponding wavelength is of the order of the cosmological scale that gravity has been tested and at the same time have the rest of the KK tower wavelengths below 0.2mm. In this scenario the gravitational interactions are due to the net effect of the massless graviton and the first ultralight KK state. A similar situation can be realized in the $''++''$ model as well. From (4.29) it can be understood that in the symmetric $''++''$ configuration the massless graviton and the special KK state contribute by the same amount to the gravitational interactions. In other words:

$$\frac{1}{M_{\rm Pl}^2}=\frac{1}{M_*^2}+\frac{1}{M_*^2}\quad\Rightarrow\quad M_{\rm Pl}=\frac{M_*}{\sqrt{2}} \tag{4.30}$$

However, if our aim is to see modifications of gravity at ultralarge distances, this is impossible because the Compton wavelength of our ultralight graviton will be e^{ky_0} times bigger than the horizon H^{-1} of the AdS_4 space on our brane due to (4.12), (4.26). The "Hubble" parameter follows m_2 rather than m_1. Moreover, in the present model, the fact that the effective four dimensional cosmological constant should be set very close to zero, requires that the warp factor is constrained by $ky_0 \geq 135$ and thus, in this case, the spectrum of the remaining KK states will be very dense (tending to continuum) bringing more states close to zero mass. The KK states that have masses which correspond to wavelengths larger than 1mm have sufficiently small coupling so that there is no conflict with phenomenology (the situation is exactly similar to the RS2 case where the coupling of the KK states is proportional to their mass and thus it is decreasing for lighter KK states). The fact that the spectrum tends to a continuum shadows the special role of the first KK state.

It is interesting to note that at the limit where the minimum of the warp factor is sent to infinity (z_∞) the special behaviour of the first KK persists and does not catch the other levels (by changing its scaling law) as was the case in $''+--+''$ model of chapter 2. This means that the limit $z\to z_\infty$ will indeed be identical to *two* RS2, but on the other hand we should emphasize that what we call graviton in the RS2 limit is actually the combination of a massless graviton *and* the "massless" limit of the special first KK state. This "massless" limit exists as we will see in the next section and ensures that locality is respected by the model, since physics on the brane does not get affected from the boundary condition at infinity.

Finally, let us note that one can build a crystalline model as in the flat brane case but without the need to introduce negative tension branes. In this case one expects again to get a band structure for the KK continuum with the width of the first band (starting from zero) much smaller than the width of the first forbidden zone. However, as in the bigravity case the width of the first band which will mediate normal four dimensional gravity at intermediate distances will be linked to the remnant cosmological constant on the branes and quite probably one will get modifications of gravity for distances much bigger than the AdS horizon. The analogue, however, of the quasi-localized model cannot be realized in the AdS brane case as it can be seen from (4.4), (4.5) since an asymptotically constant warp factor is forbidden.

2 Graviton propagator in dS_4 and AdS_4 space

In this section we will demonstrate how the van Dam-Veltman-Zakharov no-go theorem for the non-decoupling of the extra polarization states of the massive graviton can be evaded in dS_4 or AdS_4 space. For the case of dS_4 there has been a proof in [46,47], but we will present an alternative one in this section together with the case of AdS_4 for the sake of completeness. In more detail, as we have discussed at the end of chapter 2, in flat spacetime the $m \to 0$ limit of the massive graviton propagator does not give the massless one due to the non-decoupling of the additional longitudinal components. This generates the well known discontinuity between massive and massless states. Considering the massive graviton propagator in dS_4 or AdS_4 spacetime we can show that this result persists if $m/H \to \infty$ where H is the "Hubble" parameter, *i.e.* the discontinuity is still present in the $m \to 0$ limit if it happens that $m/H \to \infty$. However, in the case that $m/H \to 0$, we will explicitly show that the $m \to 0$ limit is smooth. This is an important result since it gives us the possibility to circumvent the van Dam - Veltman - Zakharov no go theorem about the non-decoupling of the extra graviton polarizations. Thus, in the limit that $m/H \to 0$ all the extra polarizations of the graviton decouple, giving an effective theory of a massless graviton with just two polarization states.

Our aim is to find the form of the massless and massive graviton propagators in the case of dS_4 and AdS_4 spacetime with arbitrary "Hubble" parameter H and graviton mass m. We wish to examine the behaviour of these propagators in the limit where $H \to 0$ and the limit where the mass of the graviton tends to zero. For simplicity we will do our calculations in Euclidean dS_4 or AdS_4 space. We can use for metric the one of the stereographic projection of the sphere or the hyperboloid[1]:

$$ds^2 = \frac{\delta_{\mu\nu}}{\left(1 \mp \frac{H^2 x^2}{4}\right)^2} dx^\mu dx^\nu \equiv g^0_{\mu\nu} dx^\mu dx^\nu \tag{4.31}$$

where $x^2 = \delta_{\mu\nu} x^\mu x^\nu$ and the scalar curvature is $R = \mp 12\ H^2$. From now on, the upper sign corresponds to AdS_4 space while the lower to dS_4 space. The fundamental invariant in these spaces is the geodesic distance $\mu(x, y)$ between two points x and y. For convenience, we will introduce another invariant u which is related with the geodesic distance by the relation $u = \cosh(H\mu) - 1$ for AdS_4 $\;(u \in [0, \infty))$ and the relation $u = \cos(H\mu) - 1$ for dS_4 $(u \in [-2, 0])$. In the small distance limit $u \sim \pm\frac{\mu^2 H^2}{2}$.

---

[1] Note that [68] and [69] whose results we use in the following have a different metric convention, but this makes no difference for our calculations.

This background metric is taken by the the Einstein-Hilbert action:

$$\mathcal{S}_G = \int d^4x\sqrt{g}\left(2M_4^2 R - \Lambda\right) \tag{4.32}$$

where the cosmological constant is $\Lambda = \mp 12\ H^2 M_4^2$ and M_4 the four dimensional fundamental scale. The spin-2 massless graviton field can be obtained by the linear metric fluctuations $ds^2 = \left(g^0_{\mu\nu} + h_{\mu\nu}\right) dx^\mu dx^\nu$. This procedure gives us the analogue of the Pauli-Fierz graviton action in curved space:

$$\begin{aligned}\frac{\mathcal{S}_0}{2M_4^2} = \int d^4x\sqrt{g^0}\Bigg\{ & -\frac{1}{4}h\Box^0 h + \frac{1}{2}h^{\mu\nu}\nabla^0_\mu\nabla^0_\nu h + \frac{1}{4}h^{\mu\nu}\Box^0 h_{\mu\nu} - \frac{1}{2}h^{\mu\nu}\nabla^0_\nu\nabla^0_\kappa h^\kappa_\mu \\ & \pm\frac{1}{2}H^2\left(\frac{1}{2}h^2 + h_{\mu\nu}h^{\mu\nu}\right)\Bigg\}\end{aligned} \tag{4.33}$$

The above action is invariant under the gauge transformation $\delta h_{\mu\nu} = \nabla^0_\mu\xi_\nu + \nabla^0_\nu\xi_\mu$ which guarantees that the graviton has only two physical degrees of freedom. This is precisely the definition of masslessness in dS_4 or AdS_4 space (for example see [70] and references therein).

The propagator of the above spin-2 massless field can be written in the form:

$$G^0_{\mu\nu;\mu'\nu'}(x,y) = (\partial_\mu\partial_{\mu'}u\partial_\nu\partial_{\nu'}u + \partial_\mu\partial_{\nu'}u\partial_\nu\partial_{\mu'}u)G^0(u) + g_{\mu\nu}g_{\mu'\nu'}E^0(u) + D[\cdots] \tag{4.34}$$

where $\partial_\mu = \frac{\partial}{\partial x^\mu}$, $\partial_{\mu'} = \frac{\partial}{\partial y^{\mu'}}$. The last term, denoted $D[\cdots]$, is a total derivative and drops out of the calculation when integrated with a conserved energy momentum tensor. Thus, all physical information is encoded in the first two terms.

The process of finding the functions G^0 and E^0 is quite complicated and is the result of solving a system of six coupled differential equations [68]. We will present here only the differential equation that G^0 satisfies to show the difference between AdS_4 and dS_4 space. This equation results from various integrations and has the general form:

$$u(u+2)G^0(u)'' + 4(u+1)G^0(u)' = C_1 + C_2 u \tag{4.35}$$

where the constants C_1 and C_2 are to be fixed by the boundary conditions. For the case of the AdS_4 space [68], these constants were set to zero so that the G^0 function vanishes at the boundary at infinity ($u \to \infty$). Using the same condition also for the

E^0 function, the exact form of them was found to be:

$$\begin{aligned} G^0(u) &= \frac{1}{8\pi^2 H^2}\left[\frac{2(u+1)}{u(u+2)} - \log\frac{u+2}{u}\right] \\ E^0(u) &= -\frac{H^2}{8\pi^2}\left[\frac{2(u+1)}{u(u+2)} + 4(u+1) - 2(u+1)^2\log\frac{u+2}{u}\right] \end{aligned} \tag{4.36}$$

For the case of the dS_4 space we iterated the procedure of [68] imposing the condition [71] that the G^0 and E^0 functions should be non-singular at the antipodal point ($u = -2$). The constants C_1 and C_2 were kept non-zero and played a crucial role in finding a consistent solution. It is straightforward to find the full expression of these functions, but we only need to know their short distance behaviour. Then with this accuracy the answer is:

$$\begin{aligned} G^0(u) &= -\frac{1}{8\pi^2 H^2}\left[\frac{1}{u} + \log(-u)\right] + \cdots \\ E^0(u) &= \frac{H^2}{8\pi^2}\left[\frac{1}{u} + 2(u+1)^2\log(-u)\right] + \cdots \end{aligned} \tag{4.37}$$

If we define $\Pi^0(u) = \frac{1}{H^4}\frac{E^0(u)}{G^0(u)}$, then for short distances ($H^2x^2 \ll 1$) where $u \to 0$ we get:

$$\begin{aligned} g_{\mu\nu}g_{\mu'\nu'} &\to \delta_{\mu\nu}\delta_{\mu'\nu'} \\ \partial_\mu\partial_{\nu'}u &\to \mp H^2\delta_{\mu\nu'} \\ G^0(u) &\to \frac{1}{4\pi^2 H^4\mu^2} \\ \Pi^0(u) &\to -1 \end{aligned} \tag{4.38}$$

and so we recover the short distance limit of the massless flat Euclidean space propagator:

$$G^0_{\mu\nu;\mu'\nu'}(x,y) = \frac{1}{4\pi^2\mu^2}(\delta_{\mu\mu'}\delta_{\nu\nu'} + \delta_{\mu\nu'}\delta_{\nu\mu'} - \delta_{\mu\nu}\delta_{\mu'\nu'}) + \cdots \tag{4.39}$$

Of course this is just as expected.

In order to describe a spin-2 massive field it is necessary to add to the above action a Pauli-Fierz mass term:

$$\frac{\mathcal{S}_m}{2M_4^2} = \frac{\mathcal{S}_0}{2M_4^2} - \frac{m^2}{4}\int d^4x\sqrt{g^0}(h_{\mu\nu}h^{\mu\nu} - h^2) \tag{4.40}$$

By adding this term we immediately lose the gauge invariance associated with the

dS_4 or AdS_4 symmetry group and the graviton acquires five degrees of freedom and becomes massive.

The propagator of this massive spin-2 field can again be written in the form:

$$G^m_{\mu\nu;\mu'\nu'}(x,y) = (\partial_\mu\partial_{\mu'}u\partial_\nu\partial_{\nu'}u + \partial_\mu\partial_{\nu'}u\partial_\nu\partial_{\mu'}u)G^m(u) + g_{\mu\nu}g_{\mu'\nu'}E^m(u) + D[\cdots] \tag{4.41}$$

The last term of the propagator in (4.41), denoted $D[\cdots]$, is again a total derivative and thus drops out of the calculation when integrated with a conserved $T_{\mu\nu}$.

At his point we should emphasize that in case of an arbitrary massive spin-2 field, the absence of gauge invariance means that there is no guarantee that the field will couple to a conserved current. However, in the context of a higher dimensional theory whose symmetry group is spontaneously broken by some choice of vacuum metric, the massive spin-2 graviton KK states couple to a conserved $T_{\mu\nu}$. One can understand this by the following example. Consider the case of the most simple KK theory, the one with one compact extra dimension. By the time we choose a vacuum metric *e.g.* $g^0_{MN} = \mathrm{diag}\,(\eta_{\mu\nu}, 1)$, the higher dimensional symmetry is broken. If we denote the graviton fluctuations around the background metric by $h_{\mu\nu}$, $h_{\mu 5}$ and h_{55}, there is still the gauge freedom:

$$\begin{aligned}\delta h_{\mu\nu} &= \partial_\mu\xi_\nu + \partial_\nu\xi_\mu \\ \delta h_{\mu 5} &= \partial_\mu\xi_5 + \partial_5\xi_\mu \\ \delta h_{55} &= 2\partial_5\xi_5\end{aligned} \tag{4.42}$$

If we Fourier decompose these fields, their n-th Fourier mode acquires a mass $m_n \propto n$ with $n = 0, 1, 2, \ldots$, but there is mixing between them. This means for example that $h^{(n)}_{\mu\nu}$ is not a massive spin-2 eigenstate *etc.*. However, we can exploit the gauge transformations (4.42) to gauge away the massive $h^{(n)}_{\mu 5}$ and $h^{(n)}_{55}$ and construct a pure spin-2 field (see for example [72] and references therein). For a comprehensive account of KK theories see [73]. The new massive spin-2 field $\rho^{(n)}_{\mu\nu}$ is invariant under (4.42) and so its Lagrangian does not exhibit a gauge invariance of the form $\delta\rho_{\mu\nu} = \partial_\mu\chi_\nu + \partial_\nu\chi_\mu$. However, since is originates from a Lagrangian that has the gauge invariance (4.42), it is bound to couple to a conserved $T_{\mu\nu}$. The argument goes on for more complicated choices of vacuum metric as for example warped metrics.

Again the functions G^m and E^m result from a complicated system of differential equations [69]. In that case, the differential equation that G^m satisfies is:

$$u(u+2)G^m(u)'' + 4(u+1)G^m(u)' \mp \left(\frac{m}{H}\right)^2 G^m(u) = C_1 + C_2 u \tag{4.43}$$

where the constants C_1 and C_2 are to be fixed by the boundary conditions. For the case of the AdS_4 space [69], these constants were set to zero so that the G^0 function vanishes at the boundary at infinity. Imposing additionally the condition of fastest falloff at infinity ($u \to \infty$) [71], the exact form of the G^m and E^0 function was found to be:

$$\begin{aligned}
G^m(u) &= \frac{\Gamma(\Delta)\Gamma(\Delta-1)}{16\pi^2\Gamma(2\Delta-2)H^2}\left(\frac{2}{u}\right)^{\Delta} F(\Delta,\Delta-1,2\Delta-2,-\frac{2}{u}) \\
E^m(u) &= -\frac{2}{3}\,\frac{\Gamma(\Delta-1)H^2}{16\pi^2\Gamma(2\Delta-2)[2+(m/H)^2]}\left(\frac{2}{u}\right)^{\Delta}\times \\
&\times \left\{ \begin{array}{l} \quad 3[2+(m/H)^2]\Gamma(\Delta-2)u^2F(\Delta-1,\Delta-2,2\Delta-2,-\frac{2}{u}) \\ -3(u+1)uF(\Delta-1,\Delta-1,2\Delta-2,-\frac{2}{u}) \\ +[3+(m/H)^2]\Gamma(\Delta)F(\Delta,\Delta-1,2\Delta-2,-\frac{2}{u}) \end{array} \right\}
\end{aligned} \tag{4.44}$$

where $\Delta = \frac{3}{2} + \frac{1}{2}\sqrt{9+4(m/H)^2}$.

For the case of the dS_4 space we iterated the procedure of [69] imposing the condition [71] that the G^m and E^m functions should be non-singular at the antipodal point ($u = -2$) and also finite as $m \to 0$. Again we kept the constants C_1 and C_2 non-zero to obtain a consistent solution. It is straightforward to find the full expression of these functions, but we only need to know their short distance behaviour. Then with this accuracy the answer is:

$$\begin{aligned}
G^m(u) &= \frac{\Gamma(\Delta)\Gamma(3-\Delta)}{16\pi^2H^2}\left[F(\Delta,3-\Delta,2,\frac{u+2}{2})-1\right]+\cdots \\
E^m(u) &= -\frac{2}{3}\,\frac{\Gamma(\Delta)\Gamma(3-\Delta)H^2}{4\pi^2[2-(m/H)^2]}\times \\
&\times \left\{ \begin{array}{l} -3[2-(m/H)^2]\left[\frac{2(u+2)}{(\Delta-1)(\Delta-2)}F(\Delta-1,2-\Delta,2,\frac{u+2}{2})+\frac{u(u+2)}{2}\right] \\ -3(u+1)\left[\frac{2}{(\Delta-1)(\Delta-2)}F(\Delta-1,2-\Delta,1,\frac{u+2}{2})+(u+1)\right] \\ +[3-(m/H)^2]\left[F(\Delta,3-\Delta,2,\frac{u+2}{2})-1\right] \end{array} \right\}+\cdots
\end{aligned} \tag{4.45}$$

where $\Delta = \frac{3}{2} + \frac{1}{2}\sqrt{9-4(m/H)^2}$.

If we define $\Pi^m(u) = \frac{1}{H^4}\frac{E^m(u)}{G^m(u)}$, then for short distances ($H^2x^2 \ll 1$) where $u \to 0$ we get:

$$G^m(u) \;\to\; \frac{1}{4\pi^2H^4\mu^2}$$

$$\Pi^m(u) \quad \rightarrow \quad -\frac{2}{3}\ \frac{3 \pm \left(\frac{m}{H}\right)^2}{2 \pm \left(\frac{m}{H}\right)^2} \tag{4.46}$$

It is interesting to consider two massless flat limits. In the first one $m \rightarrow 0$ and $H \rightarrow 0$ while $m/H \rightarrow \infty$. In this case, from (4.46) we see that we recover the Euclidean propagator for a massive graviton in flat space:

$$G^m_{\mu\nu;\mu'\nu'}(x,y) = \frac{1}{4\pi^2\mu^2}(\delta_{\mu\mu'}\delta_{\nu\nu'} + \delta_{\mu\nu'}\delta_{\nu\mu'} - \frac{2}{3}\delta_{\mu\nu}\delta_{\mu'\nu'}) + \cdots \tag{4.47}$$

This is in agreement with the van Dam - Veltman - Zakharov theorem. The second limit has $m \rightarrow 0$ and $H \rightarrow 0$ but $m/H \rightarrow 0$. In this case the propagator passes smoothly to the one of the flat massless case (4.39):

$$G^0_{\mu\nu;\mu'\nu'}(x,y) = \frac{1}{4\pi^2\mu^2}(\delta_{\mu\mu'}\delta_{\nu\nu'} + \delta_{\mu\nu'}\delta_{\nu\mu'} - \delta_{\mu\nu}\delta_{\mu'\nu'}) + \cdots \tag{4.48}$$

This is in contrary to the van Dam - Veltman - Zakharov discontinuity in flat space.

In general, we may consider the limit with m/H finite. Then for small m/H the contribution to the $\delta_{\mu\nu}\delta_{\mu'\nu'}$ structure is $-1 \pm m^2/6H^2$. Since observations agree to 0.02% accuracy with the prediction of Einstein gravitational theory for the bending of light by the sun, we obtain the limit $\frac{m}{H} \lesssim 0.04$. However, we should point out that the above results are meaningful only for the AdS case because in the dS there are no unitary spin-2 representations in the region $0 < m/H < 2$ as it was shown in [46, 47].

3 Discussion

Let us now go back to the $'' + +''$ bigravity model and see how the extra polarization states problem is evaded in the absence of a ghost radion mode. In this model we have a clear realization of the case where $m/H \rightarrow 0$. Indeed the massive graviton, which in the symmetric configuration is responsible for half of the gravitational interactions that we feel in intermediate distances, has this characteristic ratio to be $m_1/H \sim e^{-ky_0}$. This easily accommodates the bound that we presented earlier. Then, the Euclidean propagator (in configuration space) of the massive KK states for relatively large y_0 will be given by:

$$G^m_{\mu\nu;\mu'\nu'}(x,x') = \frac{1}{4\pi^2\mu^2}(\delta_{\mu\mu'}\delta_{\nu\nu'} + \delta_{\mu\nu'}\delta_{\nu\mu'} - \left(1 - \frac{1}{6}e^{-2ky_0}\right)\delta_{\mu\nu}\delta_{\mu'\nu'}) \tag{4.49}$$

where μ is the geodesic distance between two points. In the above, we have omitted terms that do not contribute when integrated with a conserved $T_{\mu\nu}$. For $ky_0 \gtrsim 2.3$ there is no problem with the bending of light since the extra polarization states are practically decoupled. Additionally, as we discussed earlier the large scale modifications of gravity are impossible to be observed because the Compton wavelength of our ultralight graviton is exponentially larger than the horizon H^{-1} of the AdS_4 space on our brane. The facts that there is no observable deviation from normal four dimensional Einsteinian gravity in this model and that there is a cosmological constant of negative sign present, make this model phenomenologically disfavoured.

Let us now make some further comments regarding the (dis)continuity of the graviton propagator. It was shown in [74, 75] that the extra polarizations are resurrected in quantum loops and thus the discontinuity reappears in the quantum level. However, purely four dimensional theory with massive gravitons where the mass of the graviton is added by hand is not well-defined as it was shown in [76]. If instead one starts from a higher dimensional theory in which the mass of the gravitons is generated dynamically, as we are doing in the models we have presented so far, and if one takes into account the whole KK graviton tower in the quantum loops, this quantum discontinuity is expected to be absent. It is worth noting that in [77] it was suggested that the extra polarization states of the massive graviton in the one AdS_4 brane model [43] can be gauged away and thus the discontinuity is absent.

Moreover, one expects that the smoothness of the limit $m \rightarrow 0$ is not only a property of the AdS_4 space but holds for any background where the characteristic curvature invariants are non-zero. For physical processes taking place in some region of a curved space with some non-zero characteristic average curvature, the effect of graviton mass is controlled by positive powers of the ratio m^2/R^2 where R^2 is a characteristic curvature invariant (made from Riemann and Ricci tensors or scalar curvature).

Finally, the conjecture that there is a smooth limit for phenomenologically observable amplitudes in brane gravity with ultralight gravitons is also supported by results based on a very interesting paper [50]. In that paper it was shown that the argument of van Dam, Veltman and Zakharov that no mass for the graviton is allowed in flat background is not entirely robust. In more detail, is was shown that there is a smooth limit for a metric around a spherically symmetric source with a mass M in a theory with massive graviton with mass m for distances smaller than a critical one which is $m^{-1}(mM/M_{\mathrm{Pl}}^2)^{1/5}$. This critical distance for a graviton of wavelength of the size of the observable universe and for a massive body of the mass of the Sun, is of the order of the solar system size. In order to see this smooth limit one should perform a non-perturbative analysis [78] since in this regime the lowest order approximation breaks down. The discontinuity reappears for distances larger than this critical distance where the linear approximation is valid. This breakdown of perturbativity, is due to the fact that the lowest order approximation calculation does not take into account the characteristic mass scale of the problem which in this case is the mass of the Sun. Since the bending of the light is observed in the first non-perturbative region, according to [78] it is impossible to rule out the massive graviton proposal.

However, the non-perturbative solution discussed in [50, 78] was found in a limited range of distance from the center and it is still unclear if it can be smoothly continued to spatial infinity (this problem was stressed in [76]). Existence of this smooth continuation depends on the full nonlinear structure of the theory. If one adds a mass term by hand the smooth asymptotic at infinity may not exit. However, it seems plausible that in all cases when modification of gravity at large distances comes from consistent higher-dimensional models, the global smooth solution can exist because in this case there is a unique non-linear structure related to the mass term which is dictated by the underlying higher-dimensional theory.

Chapter 5

Multigravity in six dimensions

As we have discussed, in five dimensions it is impossible to have flat brane multigravity models without negative tension branes. For this reason, we will consider models in six dimensions. The literature on six dimensional constructions is already quite rich [79,80, 81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97]. In this chapter we explicitly show that the difficulties of the five dimensional models can be evaded in six dimensions. It is possible to construct multigravity models with only flat positive tension branes. This is due to the fact that in six dimensions the internal space is not trivial and its curvature allows bounce configurations with the above feature. The branes which localize gravity in this setup are four-branes, but one of their dimensions is compact, unwarped and of Planck length. Thus, in the low energy limit the spacetime on the brane appears three dimensional. In order that these constructions are realized it is crucial that the tensions and/or the bulk cosmological constant are anisotropic. The five dimensional constructions in this setup come into two types. One of them involves conical singularities at finite distance from the four branes. These conical singularities support three-branes which can be of positive tension if one has an angle deficit, of zero tension if one has no angle deficit and of negative tension if one has an angle excess. The other type of the multigravity models has no conical singularity at all. It is crucial to note that there is no five dimensional effective theory for these constructions, otherwise one would get all the problems faced in the five dimensional constructions. The low energy effective theory is directly four dimensional. These constructions give for the first time a theoretically and phenomenologically viable realization of multigravity.

In the following section we present the cylindrically symmetric single four-brane warped models and show the cases where gravity is localized on the four-branes. In section two we paste two of the single four-brane solutions and obtain the double four-brane bigravity models. In section three we generalize the quasi-localized and crystalline constructions in six dimensions. Finally, we discuss the weaker energy condition in these six dimensional configurations.



Figure 5.1: The minimal single four-brane model warp factors $\sigma(\rho)$ and $\gamma(\rho)$.

1 Single brane models in six dimensions

At first we will discuss the single brane solutions in six dimensions to get an insight on the more complicated multigravity configurations. In the following subsection we will review the minimal model where the bulk cosmological constant is isotropic and then a more generalized model with arbitrary cosmological constant components along the four dimensional and the extra dimensional directions. In the following we will implicitly assume orbifolding around the four-brane.

1.1 The minimal single brane model

The simplest single brane model consists of a four-brane embedded in six dimensional *AdS* space [86]. One of the longitudinal dimensions of the four brane is compactified to a Planck length radius R while the dimensions transverse to the four-brane compact dimension are infinite. We consider the following ansatz where four dimensional Poincaré invariance is respected[1]:

$$ds^2 = \sigma(\rho)\eta_{\mu\nu}dx^\mu dx^\nu + d\rho^2 + \gamma(\rho)d\theta^2 \tag{5.1}$$

where θ is the compactified dimension with range $[0, 2\pi]$ and ρ is the infinite radial dimension.

It is straightforward to solve the Einstein equations for the bulk energy momentum tensor $T^{(B)}{}_M^{\ N} = -\Lambda\delta_M^N$ with the brane contribution:

$$T^{(br)}{}_M^{\ N} = -\delta(\rho)\begin{pmatrix} V\ \delta_\mu^\nu & & \\ & 0 & \\ & & V \end{pmatrix} \tag{5.2}$$

---

[1] Note that in this thesis we use different metric signature and different definition of the fundamental scale from [86].

From the form of the Einstein equations given in Appendix C we see that the solution for the two warp factors is:

$$\sigma(\rho) = e^{-k\rho} \quad , \quad \gamma(\rho) = R^2 e^{-k\rho} \tag{5.3}$$

with $k^2 = -\frac{\Lambda}{10M^4}$, where M is the six dimensional fundamental scale and the arbitrary integration constant R is just the radius of the four-brane (see Fig.5.1). The Einstein equations require the usual fine tuning between the bulk cosmological constant and the tension of the four brane:

$$V = -\frac{8\Lambda}{5k} \tag{5.4}$$

Let us note at this point that in [86] this fine tuning was absent because a smooth local defect was considered instead of a four-brane. In this case, the fine tuning emerges between the different components of the defect energy momentum tensor. The physics of the four-brane idealization and the one of the defect model is the same.

The four dimensional KK decomposition can be carried out as usual by considering the following graviton perturbations:

$$ds^2 = \sigma(\rho) \left[\eta_{\mu\nu} + h_{\mu\nu}(\rho, \theta, x)\right] dx^\mu dx^\nu + d\rho^2 + \gamma(\rho) d\theta^2 \tag{5.5}$$

Here, as well as throughout this chapter, we have ignored the modulus associated to the radius R of the four-brane. This will be a massless scalar and in order not to give rise to an additional long range force it should be given mass through some stabilization mechanism.

We expand the graviton perturbations in a complete set of radial eigenfunctions and Fourier angular modes:

$$h_{\mu\nu}(\rho, \theta, x) = \sum_{n,l} \phi_{(n,l)}(\rho) e^{il\theta} h_{\mu\nu}^{(n,l)}(x) \tag{5.6}$$

The differential equation for the radial wavefunctions ϕ is:

$$\phi'' - \frac{5}{2}k\phi' + \left(m^2 - \frac{l^2}{R^2}\right) e^{k\rho}\phi = 0 \tag{5.7}$$

with normalization $\int_0^\infty d\rho \sigma \sqrt{\gamma} \phi_m^* \phi_n = \delta_{mn}$. We can convert this equation to a two dimensional Schrödinger-like equation by the following redefinitions:

$$z = \frac{2}{k}\left(e^{\frac{k}{2}\rho} - 1\right) \quad , \quad \hat{\Psi} = \sigma^{3/4}\phi \tag{5.8}$$



Figure 5.2: The form of the effective potential V_{eff} for different angular quantum numbers l for the minimal single four-brane model. The δ-function in the origin is omitted.

so that

$$-\frac{1}{2\sqrt{\gamma}}\partial_z\left(\sqrt{\gamma}\partial_z\hat{\Psi}\right)+V_{eff}\hat{\Psi}=\frac{m^2}{2}\hat{\Psi} \quad , \quad V_{eff}(z)=\frac{21k^2}{32\left(\frac{kz}{2}+1\right)^2}+\frac{l^2}{2R^2}-\frac{3k}{4}\delta(z) \quad (5.9)$$

From the form of the potential (see Fig.5.2) we can easily deduce that the angular excitations spectrum will consist of continuum starting from a gap of the order $\frac{l^2}{R^2}$ and thus can be safely ignored. For the s-wave ($l=0$) there is a normalizable zero mode which is a constant in the ρ coordinate, *i.e.* $\phi_0=const$. The KK tower for the s-waves will again form a continuum but this time gapless with wavefunctions given by [86]:

$$\phi_m=N_m e^{\frac{5}{4}k\rho}\left[J_{3/2}\left(\frac{2m}{k}\right)Y_{5/2}\left(\frac{2m}{k}e^{\frac{k}{2}\rho}\right)-Y_{3/2}\left(\frac{2m}{k}\right)J_{5/2}\left(\frac{2m}{k}e^{\frac{k}{2}\rho}\right)\right] \quad (5.10)$$

The correction to Newton's law due to the s-modes can be easily calculated and one finds that it is more suppressed than the RS2 model [86, 15]. In six dimensions the correction reads:

$$\Delta V=-\frac{1}{\mathcal{O}(M_{\rm Pl}^5)}\frac{1}{r^4} \quad (5.11)$$

1.2 The generalized single brane model

We can now try to find more general single four-brane solutions by relaxing the requirement of isotropy of the bulk and brane energy momentum tensors. In contrast to the five dimensional case, this is possible in six dimensions because there are two independent functions in the metric (see Appendix C). We will consider the following anisotropic bulk energy-momentum tensor:

$$T^{(B)}{}_M{}^N = -\begin{pmatrix} \Lambda_0\ \delta_\mu^\nu & & \\ & \Lambda_\rho & \\ & & \Lambda_\theta \end{pmatrix} \tag{5.12}$$

and allow for an anisotropic brane tension of the form:

$$T^{(br)}{}_M{}^N = -\delta(\rho)\begin{pmatrix} V_0\ \delta_\mu^\nu & & \\ & 0 & \\ & & V_\theta \end{pmatrix} \tag{5.13}$$

This anisotropy can be due to different contributions to the Casimir energy in the different directions [98, 99] (see also [73]) or due to a background three-form gauge field with non-zero field strength [84].

If we define the parameter $\alpha = \frac{2\Lambda_\rho}{\Lambda_\theta}$, then the one four-brane solutions for the warp factors are:

$$\sigma(\rho) = e^{-k\rho} \quad , \quad \gamma(\rho) = R^2 e^{-\frac{k}{4}(5\alpha-6)\rho} \tag{5.14}$$



Figure 5.3: The generalized single four-brane warp factors $\sigma(\rho)$ and $\gamma(\rho)$ for different values of the parameter α.



Figure 5.4: The form of the effective potential V_{eff} for $\alpha > 2$ for different angular quantum numbers l. The δ-function in the origin is omitted.

with $k^2 = -\frac{\Lambda_\theta}{10M^4}$, where again R is the radius of the four-brane (see Fig.5.3). The flatness of the brane is achieved by the following fine tunings of the the parameters of the bulk and brane energy momentum tensors:

$$V_\theta = -\frac{8\Lambda_\theta}{5k} \quad , \quad V_0 = \frac{5\alpha+6}{16}V_\theta \quad , \quad \Lambda_0 = \frac{5\alpha^2+12}{32}\Lambda_\theta \tag{5.15}$$

Let us note that the two components of the brane tension have the same sign as long as $\alpha > -\frac{6}{5}$. For $\alpha > \frac{6}{5}$ the internal space is shrinking as in [86], for $\alpha < \frac{6}{5}$ the internal space is growing and for $\alpha = \frac{6}{5}$ the internal space is a cylinder (see Fig.5.3).

Proceeding with the usual KK decomposition we find the following equation for



Figure 5.5: The form of the effective potential V_{eff} for $\alpha < 2$ for different angular quantum numbers l. The δ-function in the origin is omitted.

the radial wavefunction ϕ:

$$\phi'' - \frac{5\alpha+10}{8}k\phi' + \left(m^2 - \frac{l^2}{R^2}e^{\frac{5}{4}(\alpha-2)k\rho}\right)e^{k\rho}\phi = 0 \tag{5.16}$$

with normalization $\int_0^\infty d\rho\sigma\sqrt{\gamma}\phi_m\phi_n = \delta_{mn}$. With the wavefunction redefinition and the coordinate change discussed in the previous subsection, we obtain the following effective potential for the two dimensional Schrödinger-like equation:

$$V_{eff}(z) = \frac{3(5\alpha+4)k^2}{64\left(\frac{kz}{2}+1\right)^2} + \frac{l^2}{2R^2}\left(\frac{kz}{2}+1\right)^{\frac{5}{2}(\alpha-2)} - \frac{3k}{4}\delta(z) \tag{5.17}$$

For the s-wave there is a normalizable zero mode when $\alpha > -\frac{2}{5}$ which is a constant in the ρ coordinate, *i.e.* $\phi_0 = const$. The KK tower for the s-waves will again form a continuum and their wavefunctions are given by:

$$\phi_m = N_m e^{\frac{\nu}{2}k\rho}\left[J_{\nu-1}\left(\frac{2m}{k}\right)Y_\nu\left(\frac{2m}{k}e^{\frac{k}{2}\rho}\right) - Y_{\nu-1}\left(\frac{2m}{k}\right)J_\nu\left(\frac{2m}{k}e^{\frac{k}{2}\rho}\right)\right] \tag{5.18}$$

where $\nu = \frac{5}{8}(\alpha+2)$. The correction to the Newton's law due to the s-modes can be easily calculated and one finds that it is negligible as long as $\alpha \gtrsim -\frac{2}{5}$:

$$\Delta V = -\frac{1}{\mathcal{O}(M_{\rm Pl}^{2\nu})}\frac{1}{r^{2\nu-1}} \tag{5.19}$$

As far as the angular excitations $l \neq 0$ are concerned, in the case where $\alpha > 2$ the potential diverges at infinity (see Fig. 5.4), so the spectrum will be discrete starting from a scale bigger than $\frac{l^2}{R^2}$ and thus can be safely ignored. On the other hand, the potential for the case $\alpha < 2$ asymptotically vanishes (see Fig. 5.5) and so the spectrum will be gapless continuum. One should examine the contribution of the angular excitations to the Newtonian potential in this case.

2 Bigravity in six dimensions

The flat single four-brane models considered in the previous sections have the characteristic that gravity is localized on the brane in the same way as in the five dimensional analogue [15]. In this section we will show how we can construct realistic multi-localization scenarios for gravity by sewing two single four-brane solutions together. We will present two explicit examples of compact double four-brane models where bigravity is realized. Firstly, we will consider a double four-brane bigravity model which in general contains an additional three-brane associated with the existence of a conical singularity. This can be done by allowing for an anisotropic four-brane tension. In the following subsection we demonstrate how one can avoid the conical singularity by also allowing for anisotropic bulk cosmological constant. In the following we will consider the orbifold geometry S^1/Z_2 in which the two four-branes lie on the fixed points.

2.1 The conifold model

We are interested in a double four-brane generalization of the minimal single four-brane model. In order to achieve this, we still assume an isotropic bulk energy momentum tensor ${T^{(B)}}_M^{\ N} = -\Lambda\delta_M^N$ but we do not impose any constraints on the four-brane tension:

$${T^{(br)}}_M^{\ N} = -\delta(\rho)\begin{pmatrix} V_0\ \delta_\mu^\nu & & \\ & 0 & \\ & & V_\theta \end{pmatrix} \tag{5.20}$$

The Einstein equations for $\sigma(\rho)$ and $\gamma(\rho)$ in this case give the following solutions for the warp factors:

$$\sigma(\rho) = \frac{\cosh^{4/5}\left[\frac{5}{4}k(\rho-\rho_0)\right]}{\cosh^{4/5}\left[\frac{5}{4}k\rho_0\right]} \quad , \quad \gamma(\rho) = R^2\frac{\cosh^{6/5}\left[\frac{5}{4}k\rho_0\right]}{\sinh^2\left[\frac{5}{4}k\rho_0\right]}\frac{\sinh^2\left[\frac{5}{4}k(\rho-\rho_0)\right]}{\cosh^{6/5}\left[\frac{5}{4}k(\rho-\rho_0)\right]} \tag{5.21}$$

with $k^2 = -\frac{\Lambda}{10M^4}$, where we have normalized $\sigma(0) = 1$ and $\gamma(0) = R^2$ (see Fig. 5.6). From the above relations it is obvious that both $\sigma(\rho)$ and $\gamma(\rho)$ have a bounce form. However, we note that $\gamma(\rho_0) = 0$, that is, $\gamma(\rho)$ is vanishing at the minimum of the warp factor $\sigma(\rho)$. This is a general characteristic of the solutions even when the branes are non-flat. From (C.3) and (C.4) we can easily show that $\gamma(\rho) = C\frac{(\sigma'(\rho))^2}{\sigma(\rho)}$ (where C is an integration constant) which implies that whenever we have a bounce in the warp factor the function $\gamma(\rho)$ will develop a zero.

In order to examine the nature of this singularity we study the form of the metric at the vicinity of the point $\rho = \rho_0$ along the lines of [100, 101]. Taking in account that in this limit we have $\sigma(\rho) \to b^2$ and $\gamma(\rho) \to \beta^2(\rho-\rho_0)^2$ the metric becomes:

$$ds^2 = \ b^2\ \eta_{\mu\nu}dx^\mu dx^\nu + d\rho^2 + \beta^2(\rho-\rho_0)^2 d\theta^2 \tag{5.22}$$

$$\text{where} \quad b^2 = \cosh^{-4/5}\left[\tfrac{5}{4}k\rho_0\right]$$

$$\text{and} \quad \beta^2 \equiv \tfrac{25k^2R^2}{16}\tfrac{\cosh^{6/5}(\frac{5}{4}k\rho_0)}{\sinh^2(\frac{5}{4}k\rho_0)}$$



Figure 5.6: The conifold bigravity model warp factors $\sigma(\rho)$ and $\gamma(\rho)$ for the symmetric configuration where $L = 2\rho_0$. Both the $\sigma(\rho)$ and $\gamma(\rho)$ have a bounce at ρ_0. However, $\gamma(\rho)$ vanishes at this point. This point in general corresponds to a conical singularity.

From the form of the metric, it is clear that for general values of the β parameter there will be a conical singularity with a corresponding deficit angle $\delta = 2\pi(1-\beta)$. The existence of this conifold singularity is connected to the presence of a 3-brane at $\rho = \rho_0$. In order to find its tension one has to carefully examine the Einstein tensor at the vicinity of the singularity. For this reason we write the metric for the internal manifold in the conformally flat form:

$$ds^2 = b^2 \eta_{\mu\nu} dx^\mu dx^\nu + f(r)(dr^2 + r^2 d\theta^2) \tag{5.23}$$

with $f(r) = r^{2(\beta-1)}$ and $\rho - \rho_0 = \beta^{-1} r^\beta$. In these coordinates it is easy to see how the three brane appears at the conifold point. The Einstein tensor can be calculated for $\rho \to \rho_0$:

$$R_{MN} - \frac{1}{2} G_{MN} R = \begin{pmatrix} \frac{\nabla^2 \log(f(r))}{2f(r)}\, b^2\ \eta_{\mu\nu} & & \\ & 0 & \\ & & 0 \end{pmatrix} \tag{5.24}$$

where ∇^2 is the flat two dimensional Laplacian. Now, given that $\nabla^2 \log(r) = 2\pi\delta^{(2)}(\mathbf{r})$

and by comparing with:

$$R_{MN}-\frac{1}{2}G_{MN}R=-\frac{V_3}{4M^4}\frac{\sqrt{-\hat{G}}}{\sqrt{-G}}\hat{G}_{\mu\nu}\delta_M^{\mu}\delta_N^{\nu}\delta(r) \tag{5.25}$$

where $\delta^{(2)}(\mathbf{r})=\frac{\delta(r)}{2\pi r}$ (see Appendix of [97]), we find that the tension of the 3-brane is:

$$V_3=4(1-\beta)M^4=\frac{2M^4}{\pi}\delta \tag{5.26}$$

Thus, if there is angle deficit $\delta>0$ ($\beta<1$) the tension of the brane is positive, whereas if there is angle excess $\delta<0$ ($\beta>1$) the tension of the brane is negative. At the critical value $\beta=0$ there is no conical singularity at all and we have a situation where two locally flat spaces touch each other at one point.

In the following we will concentrate on the symmetric case, that is, the four-branes will be considered placed at symmetric points with respect to the minimum of the warp factor ρ_0, *i.e.* $L=2\rho_0$. For the previous solution to be consistent the tensions of the four-branes for the symmetric configuration must satisfy:

$$V_\theta=-\frac{8\Lambda}{5k}\tanh\left[\frac{5}{2}k\rho_0\right] \quad , \quad V_0=\frac{3}{8}V_\theta+\frac{8\Lambda^2}{5k^2}\frac{1}{V_\theta} \tag{5.27}$$

The brane tensions for both branes are identical. Thus, the above construction consists of two positive tension four-branes placed at the end of the compact space and an intermediate three-brane due to the conifold singularity with tension depending on the parameters of the model. In the limit $\rho_0\to\infty$ we correctly obtain two identical minimal single four-brane models for the case where $\delta\to 2\pi$ ($\beta\to 0$).

The differential equation for the radial wavefunction ϕ of the graviton excitations is:

$$\phi''+2\left(\frac{\sigma'}{\sigma}+\frac{\gamma'}{4\gamma}\right)\phi'+\left(\frac{m^2}{\sigma}-\frac{l^2}{\gamma}\right)\phi=0 \tag{5.28}$$

with normalization $\int_0^{2\rho_0}d\rho\sigma\sqrt{\gamma}\phi_m\phi_n=\delta_{mn}$.

There is an obvious normalizable zero mode with $\phi_0=$ *const.* and a tower of discrete KK states. It is easy to demonstrate that for $\delta>0$ ($\beta<1$) this model has an ultralight state, leading to a theory of bigravity without negative tension branes. This follows considering the limit $\rho_0\to\infty$ when the theory describes two identical single four-brane models each of which has a massless graviton. For finite ρ_0 only their symmetric combination remains massless but their antisymmetric combination acquires an anomalously small mass, because the difference in the modulus of the



Figure 5.7: The form of the effective potential V_{eff} for different angular quantum numbers l for the conifold bigravity model in the symmetric configuration. The δ-functions in the positions of the four-branes are omitted.

wavefunctions between the massive and the massless states is significant only near ρ_0 where the wavefunctions are exponentially small[2] (see [102] for discussion of this phenomenon for fields of all spin).

We can transform this differential equation to a two dimensional Schrödinger-like one by the coordinate change $\frac{dz}{d\rho} = \sigma^{-1/2} \equiv g(z)$ and the usual wavefunction redefinition. Then the effective potential reads:

$$V_{eff} = \frac{15}{8}\left(\frac{\partial_z g}{g}\right)^2 - \frac{3}{4}\frac{\partial_z^2 g}{g} - \frac{3}{8}\frac{\partial_z g \partial_z \gamma}{g\gamma} + \frac{l^2}{2\gamma g^2} \tag{5.29}$$

We cannot write an explicit analytic formula for of the above potential in the z-coordinates because the coordinate transformation is not invertible analytically. However, since the transformation is monotonic, we can easily sketch the form of the potential (see Fig.5.7) by calculating it in the ρ coordinates. From this procedure we see that the potential for the s-wave is finite at the conical singularity ρ_0 but has a divergence for all the angular excitations with $l \neq 0$. This means that only the s-wave excitations with communicate the two parts of the conifold and the other excitations will be confined in the two semicones. The spectrum of KK states in this case will be discrete.

---

[2] There is, however, a mismatch of the degrees of freedom of the massive and massless graviton, which may not be phenomenologically dangerous, but still raises the issue of locality in the infinite four-brane separation limit.

The construction may be readily modified to obtain a six-dimensional analogue of the locally localized model [43]. This is done by considering the asymmetric situation in which one of the fixed point four-branes is moved to infinity (in the asymmetric case the tensions of the two four-branes are different). Then gravity on the four-brane at $\rho = 0$ will be mediated by only the ultralight state since the graviton zero mode will not be normalizable.

2.2 The non-singular model

The appearance of a conifold singularity can be avoided if instead of using the minimal single four-brane solution to build the double four-brane solution, one uses the generalized single four-brane model. By allowing for an isotropic bulk cosmological constant one can arrange that the function $\gamma(\rho)$ does not develop a zero. In this case the (θ, θ) and (ρ, ρ) part of Einstein equations in the bulk, (C.3) and (C.4), have solutions:

$$\sigma(\rho) = \frac{\cosh^{4/5}\left[\frac{5}{4}k(\rho-\rho_0)\right]}{\cosh^{4/5}\left[\frac{5}{4}k\rho_0\right]} \quad , \quad \gamma(\rho) = A\frac{\sinh^{\alpha}\left[\frac{5}{4}k(\rho-\rho_0)\right]}{\cosh^{6/5}\left[\frac{5}{4}k(\rho-\rho_0)\right]} \tag{5.30}$$

where A is a constant and $\alpha \equiv \frac{2\Lambda_\rho}{\Lambda_\theta}$. The (μ, ν) component of the Einstein equations (C.5), however, restricts the possible values of α to $\alpha = 2$ or $\alpha = 0$. The case of $\alpha = 2$ corresponds to the previous conifold model. For the case $\alpha = 0$, which corresponds to the choice $\Lambda_\rho = 0$, the solution is:

$$\sigma(\rho) = \frac{\cosh^{4/5}\left[\frac{5}{4}k(\rho-\rho_0)\right]}{\cosh^{4/5}\left[\frac{5}{4}k\rho_0\right]} \quad , \quad \gamma(\rho) = R^2\frac{\cosh^{6/5}\left[\frac{5}{4}k\rho_0\right]}{\cosh^{6/5}\left[\frac{5}{4}k(\rho-\rho_0)\right]} \tag{5.31}$$

with $k^2 = -\frac{\Lambda_\theta}{10M^4}$, where we have normalized $\sigma(0) = 1$ and $\gamma(0) = R^2$ (see Fig. 5.8). From the above the absence of any singularity is obvious since $\gamma(\rho)$ does not vanish at any finite value of ρ.

The four branes will appear as δ-function singularities at the points where we will cut the space in the ρ direction. In the following we will concentrate on the symmetric case, that is, the branes will be considered placed in symmetric points in respect to the extremum of the warp factor ρ_0, *i.e.* $L = 2\rho_0$. In order for the above configuration to be realized, the components of the four-brane tensions and the bulk cosmological constant must be tuned to give:

$$V_\theta = -\frac{8\Lambda}{5k}\tanh\left[\frac{5}{2}k\rho_0\right] \quad , \quad V_0 = \frac{3}{8}V_\theta \quad , \quad \Lambda_0 = \frac{3}{8}\Lambda_\theta \tag{5.32}$$



Figure 5.8: The non-singular bigravity model warp factors $\sigma(\rho)$ and $\gamma(\rho)$ for the symmetric configuration where $L = 2\rho_0$. The $\sigma(\rho)$ warp factor has exactly the same form as in the conifold model. The $\gamma(\rho)$ warp factor however now has an inverse bounce form. In this case $\gamma(\rho)$ does not vanish anywhere and thus the model is free of singularities.

The brane tensions for both branes are identical. Thus, we have constructed a compact model with two positive tension flat branes with a bounce in the warp factors. Note the extra condition relating the different components of the bulk cosmological constant; this anisotropy in the bulk cosmological constant is what allows us to have a non-singular solution. In the limit $\rho_0 \to \infty$ we obtain two identical generalized single four-brane $\alpha = 0$ models.

The differential equation for the radial wavefunction ϕ of the graviton excitations is given by (5.28) for the relevant $\sigma(\rho)$ and $\gamma(\rho)$ functions. Although it is difficult to find an analytical solution of the corresponding differential equation for the KK states, the existence of a light special KK is assured from the locality arguments discussed above. Since at the infinite brane separation limit we recover two identical generalized single four-brane models (for $\alpha = 0$) where each of them supports a massless zero mode, at a



Figure 5.9: The form of the effective potential V_{eff} for different angular quantum numbers l for the non-singular bigravity model in the symmetric configuration. The δ-functions in the positions of the four-branes are omitted.

finite separation configuration the one of the zero modes will become the special light KK state.

By making an appropriate change of coordinates and a redefinition of the wavefunction we can bring the previous differential equation in a two dimensional Schrödinger form with the effective potential (5.29) that is plotted in Fig.5.9. Note that in this case the angular excitations feel the whole region between the two positive branes and thus communicate physics between them. The spectrum of KK states in this case will again be discrete.

3 Multigravity in six dimensions

In this section we will show how we can obtain the analogues of the GRS model of quasi-localized gravity and of the crystal universe in six dimensions, in both cases without the need of introducing moving negative tension branes.

3.1 Quasi-localized gravity

As in the bigravity models we have two possible ways of realizing a quasi-localized scenario in six dimensions. The first model can be built from the $\alpha = 0$ bigravity model, by cutting the space at a point ρ_b on the right of the position of the bounce and sewing it to flat space on the right (see Fig.5.10). In that case we will have a system



Figure 5.10: The $\alpha = 0$ quasi-localized model warp factors $\sigma(\rho)$ and $\gamma(\rho)$. The bulk region for $\rho > \rho_b$ is flat.

of two positive tension branes and by appropriately tuning the different components of the brane tension of the second brane we can find solutions where the warp factor is constant in the flat bulk region. The fine tunings which achieve this are:

$$V_\theta^{(2)} = -\frac{4\Lambda}{5k}\tanh\left[\frac{5}{2}k(\rho_b - \rho_0)\right] \quad , \quad V_0^{(2)} = \frac{3}{8}V_\theta^{(2)} \tag{5.33}$$

From the above formulas it is obvious that if we paste the flat space at the position of the bounce, $i.e$ $\rho_b = \rho_0$ then the second brane is tensionless. If we relax these fine tunings we can find solutions where the warp factors in the flat bulk region are non trivial, but we will not discuss this possibility here.

The above construction will obviously have no normalizable zero mode due to the infinite volume of the system and the KK spectrum will be continuous. However, as in the five dimensional case we expect that the low part of the KK tower will have a “resonance”-like coupling to matter on the $''+''$ brane at $\rho = 0$. This will mediate normal gravity at intermediate distances but will change the nature of the gravitational law at ultralarge scales.



Figure 5.11: The $\alpha = 2$ quasi-localized model warp factors $\sigma(\rho)$ and $\gamma(\rho)$. The bulk region for $\rho > \rho_b$ is flat.

The second model of quasi-localized gravity can be built in a similar fashion from the $\alpha = 2$ bigravity model (see Fig.5.11). Again we obtain a system of two positive tension branes which will have a constant warp factor in the flat bulk region when the following fine tunings are demanded:

$$V_\theta^{(2)} = -\frac{4\Lambda}{5k}\tanh\left[\frac{5}{2}k(\rho_b - \rho_0)\right] \quad , \quad V_0^{(2)} = \frac{3}{8}V_\theta^{(2)} + \frac{8\Lambda^2}{5k^2}\frac{1}{V_\theta^{(2)}} \tag{5.34}$$

From the above formulas it is obvious that if we paste the flat space at the position of the bounce, *i.e* $\rho_b = \rho_0$ then the tension $V_\theta^{(2)}$ vanishes whereas the tension $V_0^{(2)}$ diverges. This limit is not well behaved since the internal space becomes singular in a whole line. At this limit obviously classical gravity breaks down and one would expect that quantum gravity corrections would resolve the singularity. However, one should check if these quantum gravity corrections affect the solution in the vicinity of the four-brane at $\rho = 0$, *e.g.* by examining the R^2 correction to the Einstein action.

The above construction will again have no normalizable zero mode and the KK spectrum will be continuous. A "resonance"-like coupling of the KK states to matter on the $''+''$ brane at $\rho = 0$ is again expected which will give normal four dimensional



Figure 5.12: The $\alpha = 0$ crystalline model warp factors $\sigma(\rho)$ and $\gamma(\rho)$.

gravity at intermediate distances but will change the nature of the gravitational law at ultralarge scales.

3.2 The crystal universe model

Another obvious generalization of the five dimensional multigravity models is the one of the crystalline brane model. All that one has to do is to paste an infinite array of bigravity models. We again take this constructions in two copies. One for $\alpha = 0$ (Fig.5.12) and one for $\alpha = 2$ (Fig.5.13). A band structure is again expected as in the five dimensional case and the width of the first band will be exponentially smaller than the one of the first forbidden zone. As far as the phenomenology is concerned, these models will generate normal gravity at intermediate scales as the first band will behave as an effective zero mode, whereas at ultralarge scales will have the same change of the gravitational law as the one of the corresponding quasi-localized models.



Figure 5.13: The $\alpha = 2$ crystalline model warp factors $\sigma(\rho)$ and $\gamma(\rho)$.

4 The weaker energy condition

In this chapter we have constructed, for the first time, flat brane theories which can lead to multigravity models and their associated modifications of gravity at large distances *without* introducing moving negative tension branes. The constructions are made possible by going to a six dimensional theory. In five dimensions with flat branes, the presence of a bounce of the warp factor was linked to the violation of the weaker energy condition. This is not true, however, in six dimensions as we will show in this section. In five dimensions with the metric:

$$ds^2 = e^{-A(\rho)}\eta_{\mu\nu}dx^\mu dx^\nu + d\rho^2 \tag{5.35}$$

one can readily show that the weaker energy condition requires that:

$$A'' \geq 0 \tag{5.36}$$

which is violated at the position of the moving negative tension branes. In the case that the branes were AdS one could have a bounce (without having negative tension branes) and still satisfy the weaker energy condition because the above relation is modified to [19]:

$$A'' \geq -2H^2 e^A \tag{5.37}$$

However, such models do not lead to modifications of gravity at large distances and moreover the remnant negative cosmological constant is in conflict with current observations.

In the six dimensional case with metric:

$$ds^2 = e^{-A(\rho)}\eta_{\mu\nu}dx^\mu dx^\nu + d\rho^2 + e^{-B(\rho)}d\theta^2 \tag{5.38}$$

from the weaker energy condition one finds two inequalities which can be cast into the following relation:

$$\frac{1}{6}(B')^2 - \frac{1}{3}B'' - \frac{1}{6}A'B' \leq A'' \leq -\frac{1}{2}(B')^2 + B'' - \frac{3}{2}A'B' + 2(A')^2 \tag{5.39}$$

These inequalities hold for the $\alpha = 2$ models as long as the three-branes sitting on the conical singularities have positive tension. It is violated, however, in the $\alpha = 0$ models everywhere in the bulk. This shows that in six dimensional flat brane models, the presence of bounces of the warp factor is not necessarily linked to whether the weaker energy condition is satisfied or not. It is not yet clear if the violation of the weaker energy condition in the $\alpha = 0$ models is a sign of a possible instability.

Chapter 6

Conclusions

In thesis we have presented a comprehensive account of the multigravity scenario and how it can be realized in warped brane-world models in five and six dimensions. It is clear that the possibility of having massive gravitons contributing significantly to intermediate distance gravitational interactions is not excluded by current observations and furthermore it offers phenomenological signatures which are in principle testable. Moreover, taking into account that massive gravitons necessarily arise as KK states from the dimensional reduction of a higher dimensional theory, these modifications of gravity can be a very unusual window to extra dimensions. Multigravity models which involve brane configurations with interbrane distances of the order of the Planck length, modify gravity not only at distances of the order of the Planck length as usually do ordinary KK theories, but also at ultralarge distances. The crossover scale of normal to modified gravity is related to the interbrane distances and in that sense is generated by the geometry itself.

The first attempts to build such a model faced inconsistency problems. The $''+-+''$ bigravity model, the quasi-localized GRS model and the crystalline model, although they had promising phenomenology, they all shared the characteristic of moving negative tension branes. As we explicitly discussed in this thesis the moduli corresponding to the fluctuations of these branes are ghost fields and therefore unacceptable. However, the general characteristics of these models persist in more involved constructions that have less or no problems. For example, the appearance of light states whenever we have more than one positive branes that localize gravitons is a generic characteristic which persists in the AdS brane models or the six dimensional ones.

The AdS models on the other hand, although they do not have the ghost radion problem, do not give some interesting phenomenology. The first reason is the very fact that the remnant cosmological constant is of negative sign, in contradiction with observations. The second reason is that they do not lead to any observable modifications of gravity at any observable scale. The modifications of gravity at ultralarge scales are hidden behind the AdS horizon, while the modifications at all scales due to the different propagator structure of the massive gravitons are enormously suppressed. It is, however, interesting how the AdS brane models escape the van Dam-Veltman-Zakharov no-go theorem about the non-decoupling of the additional polarization states of the massive graviton. This in addition to the observation of Vainshtein that the latter no-go theorem may not be valid even for flat background, suggests that the massive graviton proposal does not face any fundamental obstructions.

Certainly the only theoretically and phenomenologically viable possibility of multigravity at the moment is the one in six dimensional models. Due to the non-trivial curvature of the internal space one can have bounces of the warp factor with only positive tension flat branes. There is no five dimensional effective regime in these models, since otherwise one would face all the problems present in five dimensions. The low energy theory is directly four dimensional. Moreover, the replication of the multigravity models in two types each, one with conical singularities and one without them, gives a wealth of possibilities that offer promising phenomenology.

There are a lot of open questions regarding multigravity at the moment. The most important one is the construction of a cosmological model of multigravity. So far the models we considered involved either flat or AdS branes which are not valid descriptions of our Universe at large scales where the modifications of gravity due to the massive gravitons are expected to appear. A cosmological generalization of these models would probably reveal corrections to the four dimensional Friedman equation at cosmological scales. This information is essential if one wishes to put these models into observational test.

Furthermore, as far as model building is concerned, there are a lot of issues to be investigated in the six dimensional models. A careful treatment of the angular excitations in the case of the non-singular models should be carried out to examine the nature of gravity on the four-branes. Furthermore, one should examine the moduli of the system, the radions and the dilaton, and calculate their mass. It is quite probable that the modulus corresponding to ρ_0 is massive as in the five dimensional AdS_4 branes case and so the system is self-stabilized. Finally, the non-trivial way that locality is preserved by these model should be carefully studied.

Appendix A

Effective action for graviton-radion-dilaton system

It is convenient to define a new variable z defined by:

$$\frac{1}{a(y)} = \frac{dz}{dy} \tag{A.1}$$

In the coordinates (x, z) the metric (3.10) is conformal to a flat perturbed metric $\tilde{G}_{AB} = \eta_{AB} + \tilde{H}_{AB}$:

$$\begin{aligned} ds^2 &= a^2 \left[\tilde{G}_{\mu\nu}\, dx^\mu dx^\nu + \tilde{G}_{zz}\, dz^2\right] = a^2 \left[\eta_{MN} + \tilde{H}_{MN}\, dx^M dx^N\right] \\ \tilde{H}_{\mu\nu} &= \varphi_1(x,z)\, \eta_{\mu\nu} + 2\epsilon(z)\, \partial_\mu \partial_\nu f_2(x) + \tilde{h}_{\mu\nu}(x,z) \\ \tilde{H}_{zz} &= \varphi_2(x,z) \end{aligned} \tag{A.2}$$

where the fields φ_1 and φ_2 are:

$$\begin{aligned} \varphi_1(x,z) &= Q(z) f_1(x) + B(z) f_2(x) \\ \varphi_2(x,z) &= q(z) f_1(x) + A(z) f_2(x) \end{aligned} \tag{A.3}$$

Inserting (A.2) in (3.1) and taking into account the equation of motion satisfied by a one gets:

$$\begin{aligned} &\mathcal{S}_{eff} = \int d^4x\, \mathcal{L}_{eff} = \int d^4x\, 2M^3 \int dz \left[a^3 \mathcal{L}^{(0)}_{PF}(\tilde{h}) + \frac{a^3}{4}\left[(\partial_z \tilde{h})^2 - \partial_z \tilde{h}_{\mu\nu}\, \partial_z \tilde{h}^{\mu\nu}\right] + \mathcal{L}_\varphi + \mathcal{L}_{\tilde{h}\varphi}\right] \\ &\quad \mathcal{L}_\varphi = \mathcal{L}_1 + \mathcal{L}_2 + \mathcal{L}_{12} \end{aligned} \tag{A.4}$$

where

$$\begin{aligned}\mathcal{L}_{\tilde{h}\varphi} &= \left[a^3\left(\varphi_1 + \frac{1}{2}\varphi_2\right) + f_2\frac{d}{dz}\left(\epsilon' a^3\right)\right]\partial_\mu\partial_\nu\tilde{h}^{\mu\nu} \\ &\quad - \left[a^3\left(\varphi_1 + \frac{1}{2}\varphi_2\right) + f_2\frac{d}{dz}\left(\epsilon' a^3\right)\right]\Box\tilde{h} + \frac{3}{2}\frac{d}{dz}\left(a^2a'\varphi_2 - a^3\varphi_1'\right)\tilde{h}\end{aligned} \tag{A.5}$$

$$\mathcal{L}_1 = \partial_\mu f_1\partial^\mu f_1\,\frac{3}{2}a^3\left(Q^2 + Qq\right) + f_1^2\left(3a^3Q'^2 + 3a'^2aq^2 - 6a^2a'Q'q\right) \tag{A.6}$$

$$\begin{aligned}\mathcal{L}_2 &= \partial_\mu f_2\partial^\mu f_2\left(\frac{3}{2}a^3B^2 + \frac{3}{2}a^3AB - 3a^3B'\epsilon' + 3a^2a'\epsilon' A\right) \\ &\quad + f_2^2\left(3a^3B'^2 + 3a'^2aA^2 - 6a^2a'AB'\right)\end{aligned} \tag{A.7}$$

$$\begin{aligned}\mathcal{L}_{12} &= \partial_\mu f_1\partial^\mu f_2\left(\frac{3}{2}a^3Bq + \frac{3}{2}a^3AQ + 3a^3BQ - 3a^3\epsilon' Q' + 3a^2a'\epsilon' q\right) \\ &\quad + f_1f_2\left[6a^3B'Q' + 6a'^2aAq - 6a^2a'\left(AQ' + qB'\right)\right]\end{aligned} \tag{A.8}$$

and $\mathcal{L}_{PF}^{(0)}(\tilde{h})$ is the four dimensional Pauli-Fierz Lagrangian for $\tilde{h}$ without the mass term:

$$\mathcal{L}_{PF}^{(0)}(\tilde{h}) = \frac{1}{2}\partial_\nu\tilde{h}_{\mu\alpha}\,\partial^\alpha\tilde{h}^{\mu\nu} - \frac{1}{4}\partial_\mu\tilde{h}_{\alpha\beta}\,\partial^\mu\tilde{h}^{\alpha\beta} - \frac{1}{2}\partial_\alpha\tilde{h}\,\partial_\beta\tilde{h}^{\alpha\beta} + \frac{1}{4}\partial_\alpha\tilde{h}\,\partial^\alpha\tilde{h} \tag{A.9}$$

All indices in the above expressions are raised and lowered by $\eta_{\mu\nu}$. Differentiation with respect of z is denoted with a prime. The absence of mixing terms in $\mathcal{L}_{eff}$ yields the following constraints:

$$A(z) = \frac{aB'}{a'}\,, \qquad \frac{d}{dz}\left(Ba^2\right) + \frac{2a'}{a^2}\frac{d}{dz}\left(a^3\epsilon'\right) = 0 \tag{A.10}$$

$$Q(z) = c\,a^{-2}\,, \qquad q(z) = -2c\,a^{-2}, \quad c \text{ is a constant} \tag{A.11}$$

$$\int dz\,aA(z) = 0 \tag{A.12}$$

Equations (A.10)-(A.12) give:

$$\begin{aligned}\mathcal{L}_{eff} = 2M^3 \int dz \Big\{ & a^3 \mathcal{L}^{(0)}_{PF}(\tilde{h}) + \frac{a^3}{4}\left[(\partial_z \tilde{h})^2 - \partial_z \tilde{h}_{\mu\nu}\, \partial_z \tilde{h}^{\mu\nu}\right] \\ & - \frac{3}{2} c^2 a^{-1} \partial_\mu f_1 \partial^\mu f_1 + \frac{3}{4}\frac{a^2}{a'}\frac{d}{dz}\left(B^2 a^2\right) \partial_\mu f_2 \partial^\mu f_2 \Big\}\end{aligned} \tag{A.13}$$

In particular the effective Lagrangian $\mathcal{L}_{Scal}$ for the dilaton f_1 and the radion f_2 is:

$$\begin{aligned}\mathcal{L}_{Scal} &= \mathcal{K}_1\, f_1 \Box f_1 + \mathcal{K}_2\, f_2 \Box f_2 \\ \mathcal{K}_1 &= 2M^3\, \frac{3}{2} c^2 \int_{-L}^{L} a^{-2}\, dy \\ \mathcal{K}_2 &= -2M^3\, \frac{3}{4} \int_{-L}^{L} a \left(\frac{da}{dy}\right)^{-1} \frac{d}{dy}\left(B^2 a^2\right)\, dy\end{aligned} \tag{A.14}$$

with:

$$\begin{aligned}&\frac{d}{dy}\left(B a^2\right) + 2a^{-1}\frac{da}{dy}\frac{d}{dy}\left(a^4 \partial_y \epsilon\right) = 0 \\ &\int_{-L}^{L} dy\; a \left(\frac{da}{dy}\right)^{-1} \frac{dB}{dy} = 0\end{aligned} \tag{A.15}$$

As a result only $\partial_y \epsilon(0)$, $\partial_y \epsilon(r)$ and $\partial_y \epsilon(L)$ enter the radion effective action.

Appendix B

Ricci tensor and scalar for the dilaton perturbation

In this Appendix we list the Ricci tensor components, the Ricci scalar and the action obtained by the metric:

$$ds^2 = e^{-a(y)^{-2}\gamma(x)} a(y)^2 g_{\mu\nu}(x) dx^\mu dx^\nu + \left(1 + a(y)^{-2}\gamma(x)\right)^2 dy^2 \tag{B.1}$$

The spacetime components of the five dimensional Ricci tensor are:

$$\begin{aligned} R_{\mu\nu} \ = \ R_{g\ \mu\nu} \ &+ \ \frac{g_{\mu\nu}}{2} a^{-2}\Box\gamma - \frac{g_{\mu\nu}}{2}\frac{a^{-6}\gamma}{(1+a^{-2}\gamma)}\gamma_{,\kappa}\gamma^{,\kappa} - \frac{1}{2}\left(\frac{1-a^{-2}\gamma}{1+a^{-2}\gamma}\right) a^{-4}\gamma_{,\mu}\gamma_{,\nu} \\ &+ \ \frac{a^{-4}\gamma}{(1+a^{-2}\gamma)} D_\mu \partial_\nu \gamma - g_{\mu\nu} e^{-a^{-2}\gamma} (a')^2 \left(\frac{3+4a^{-2}\gamma}{1+a^{-2}\gamma}\right) \\ &- \ g_{\mu\nu} e^{-a^{-2}\gamma} \frac{aa''}{(1+a^{-2}\gamma)} \end{aligned} \tag{B.2}$$

and the (55) component:

$$\begin{aligned} R_{55} \ = \ & e^{a^{-2}\gamma}(1+a^{-2}\gamma)a^{-6}\gamma_{,\mu}\gamma^{,\mu} - e^{a^{-2}\gamma}(1+a^{-2}\gamma)a^{-4}\Box\gamma \\ & -4(1+a^{-2}\gamma)a^{-1}a'' - 4(1+a^{-2}\gamma)a^{-4}(a')^2\gamma \end{aligned} \tag{B.3}$$

Finally the Ricci scalar is:

$$\begin{aligned} R \ = \ & e^{a^{-2}\gamma} a^{-2} R_g + e^{a^{-2}\gamma}\left(\frac{1+3a^{-2}\gamma}{1+a^{-2}\gamma}\right) a^{-4}\Box\gamma + \frac{1}{2} e^{a^{-2}\gamma}\left(\frac{1-3a^{-2}\gamma}{1+a-2\gamma}\right) a^{-6}\gamma_{,\mu}\gamma^{,\mu} \\ & -8\frac{a^{-1}a''}{(1+a^{-2}\gamma)} - \frac{12a^{-2}(a')^2 + 20a^{-4}(a')^2\gamma}{(1+a^{-2}\gamma)} \end{aligned} \tag{B.4}$$

In the above expressions the indices are raised and lowered by $g_{\mu\nu}$.

The action (3.48) then becomes:

$$\begin{aligned}
\mathcal{S} \;=\; & \int d^4x dy \sqrt{-g} \Bigg\{ 2M^3 \Bigg[e^{-a^{-2}\gamma} a^2 (1 + a^{-2}\gamma) R_g + e^{-a^{-2}\gamma} \left(1 + \frac{5}{2} a^{-2}\gamma + \frac{3}{2} a^{-4}\gamma^2 \right) g^{\mu\nu} D_\mu \partial_\nu \gamma \\
& \qquad + e^{-2a^{-2}\gamma} (-8a'' a^3 - 12(a')^2 a^2 - 20(a')^2 \gamma) \Bigg] \\
& \quad - \frac{1}{2} \frac{a^4}{(1 + a^{-2}\gamma)} e^{-2a^{-2}\gamma} \Phi' \cdot \Phi' - e^{-2a^{-2}\gamma} a^4 (1 + a^{-2}\gamma) V(\Phi) \\
& \quad - e^{-2a^{-2}\gamma} a^4 \sum_i \lambda_i(\Phi) \delta_{(} y - y_i) \Bigg\} \qquad \text{(B.5)}
\end{aligned}$$

Appendix C

Equations of motion in six dimensions

In this Appendix we provide the Einstein equations for the class of models that we considered in chapter 5 where the metric is:

$$ds^2 = \sigma(\rho)\eta_{\mu\nu}dx^\mu dx^\nu + d\rho^2 + \gamma(\rho)d\theta^2 \tag{C.1}$$

Ignoring the three-branes located on the conical singularities, the Einstein equations are generally written as:

$$G_{MN} = \frac{1}{4M^4}(T^{(B)}_{MN} + T^{(br)}_{MN}) \tag{C.2}$$

where $T^{(B)}_{MN}$ is the bulk energy momentum tensor of the general form (5.12) and $T^{(br)}_{MN}$ the four-brane energy momentum tensor of the form (5.13).

In the absence of a four dimensional cosmological constant, the (θ, θ) component of the above equation is:

$$2\frac{\sigma''}{\sigma} + \frac{1}{2}\left(\frac{\sigma'}{\sigma}\right)^2 = -\frac{\Lambda_\theta}{4M^4} - \frac{V^i_\theta}{4M^4}\delta(\rho - \rho_i) \tag{C.3}$$

The (ρ, ρ) component is:

$$\frac{3}{2}\left(\frac{\sigma'}{\sigma}\right)^2 + \frac{\sigma'\gamma'}{\sigma\gamma} = -\frac{\Lambda_\rho}{4M^4} \tag{C.4}$$

Finally, the (μ, ν) component is:

$$\frac{3}{2}\frac{\sigma''}{\sigma} + \frac{3}{4}\frac{\sigma'\gamma'}{\sigma\gamma} - \frac{1}{4}\left(\frac{\gamma'}{\gamma}\right)^2 + \frac{1}{2}\frac{\gamma''}{\gamma} = -\frac{\Lambda_0}{4M^4} - \frac{V^i_0}{4M^4}\delta(\rho - \rho_i) \tag{C.5}$$

These equations may be compared with the ones of the five dimensional case where the metric:

$$ds^2 = \sigma(\rho)\eta_{\mu\nu}dx^\mu dx^\nu + d\rho^2 \tag{C.6}$$

gives rise to the (ρ, ρ) component:

$$\frac{3}{2}\left(\frac{\sigma'}{\sigma}\right)^2 = -\frac{\Lambda}{4M^3} \tag{C.7}$$

and the (μ, ν) component:

$$\frac{3}{2}\frac{\sigma''}{\sigma} = -\frac{\Lambda}{4M^3} - \frac{V^i}{4M^3}\delta(\rho - \rho_i) \tag{C.8}$$

The extra freedom that we have in the six dimensional case is apparent.

Bibliography

[1] K. Akama, Lect. Notes Phys. **176**, 267 (1982) [hep-th/0001113].

[2] V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B **125**, 136 (1983).

[3] M. Visser, Phys. Lett. B **159**, 22 (1985) [hep-th/9910093].

[4] E. J. Squires, Phys. Lett. B **167**, 286 (1986).

[5] J. Polchinski, hep-th/9611050.

[6] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B **429**, 263 (1998) [hep-ph/9803315].

[7] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B **436**, 257 (1998) [hep-ph/9804398].

[8] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Rev. D **59**, 086004 (1999) [hep-ph/9807344].

[9] P. Horava and E. Witten, Nucl. Phys. B **460**, 506 (1996) [hep-th/9510209].

[10] E. Witten, Nucl. Phys. B **471**, 135 (1996) [hep-th/9602070].

[11] P. Horava and E. Witten, Nucl. Phys. B **475**, 94 (1996) [hep-th/9603142].

[12] A. Lukas, B. A. Ovrut, K. S. Stelle and D. Waldram, Phys. Rev. D **59**, 086001 (1999) [hep-th/9803235].

[13] A. Lukas, B. A. Ovrut and D. Waldram, Phys. Rev. D **60**, 086001 (1999) [hep-th/9806022].

[14] M. Gogberashvili, Mod. Phys. Lett. A **14**, 2025 (1999) [hep-ph/9904383].

[15] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999) [hep-th/9906064].

[16] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [hep-ph/9905221].

[17] P. Binetruy and J. Silk, Phys. Rev. Lett. **87**, 031102 (2001) [astro-ph/0007452].

[18] M. Bastero-Gil and L. Mersini, Phys. Rev. D **65**, 023502 (2002) [astro-ph/0107256].

[19] J. P. Uzan and F. Bernardeau, Phys. Rev. D **64**, 083004 (2001) [hep-ph/0012011].

[20] C. M. Will, Phys. Rev. D **57**, 2061 (1998) [arXiv:gr-qc/9709011].

[21] I. I. Kogan, S. Mouslopoulos, A. Papazoglou, G. G. Ross and J. Santiago, Nucl. Phys. B **584**, 313 (2000) [hep-ph/9912552].

[22] R. Gregory, V. A. Rubakov and S. M. Sibiryakov, Phys. Rev. Lett. **84**, 5928 (2000) [hep-th/0002072].

[23] C. Csaki, J. Erlich and T. J. Hollowood, Phys. Rev. Lett. **84**, 5932 (2000) [hep-th/0002161].

[24] G. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B **484**, 112 (2000) [hep-th/0002190].

[25] I. I. Kogan and G. G. Ross, Phys. Lett. B **485**, 255 (2000) [hep-th/0003074].

[26] I. I. Kogan, S. Mouslopoulos, A. Papazoglou and G. G. Ross, Nucl. Phys. B **595**, 225 (2001) [hep-th/0006030].

[27] G. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B **485**, 208 (2000) [hep-th/0005016].

[28] G. Dvali and G. Gabadadze, Phys. Rev. D **63**, 065007 (2001) [hep-th/0008054].

[29] H. van Dam and M. Veltman, Nucl. Phys. B **22**, 397 (1970).

[30] V.I. Zakharov, JETP Lett. **12** (1970) 312.

[31] G. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B **484**, 129 (2000) [hep-th/0003054].

[32] L. Pilo, R. Rattazzi and A. Zaffaroni, JHEP **0007**, 056 (2000) [hep-th/0004028].

[33] I. I. Kogan, S. Mouslopoulos, A. Papazoglou and L. Pilo, hep-th/0105255.

[34] C. Charmousis, R. Gregory and V. A. Rubakov, Phys. Rev. D **62**, 067505 (2000) [hep-th/9912160].

[35] P. Binetruy, C. Deffayet and D. Langlois, hep-th/0101234.

[36] C. Csaki, J. Erlich and T. J. Hollowood, Phys. Lett. B **481**, 107 (2000) [hep-th/0003020].

[37] R. Gregory, V. A. Rubakov and S. M. Sibiryakov, Phys. Lett. B **489**, 203 (2000) [hep-th/0003045].

[38] D. Z. Freedman, S. S. Gubser, K. Pilch and N. P. Warner, Adv. Theor. Math. Phys. **3**, 363 (1999) [hep-th/9904017].

[39] E. Witten, hep-ph/0002297.

[40] A. Papazoglou, Phys. Lett. B **505**, 231 (2001) [hep-th/0102015].

[41] G. Gibbons, R. Kallosh and A. Linde, JHEP **0101**, 022 (2001) [hep-th/0011225].

[42] I. I. Kogan, S. Mouslopoulos and A. Papazoglou, Phys. Lett. B **501**, 140 (2001) [hep-th/0011141].

[43] A. Karch and L. Randall, Int. J. Mod. Phys. A **16**, 780 (2001) [hep-th/0011156].

[44] M. D. Schwartz, Phys. Lett. B **502**, 223 (2001) [hep-th/0011177].

[45] A. Karch and L. Randall, hep-th/0105108.

[46] A. Higuchi, Nucl. Phys. B **282**, 397 (1987).

[47] A. Higuchi, Nucl. Phys. B **325**, 745 (1989).

[48] I. I. Kogan, S. Mouslopoulos and A. Papazoglou, Phys. Lett. B **503**, 173 (2001) [hep-th/0011138].

[49] M. Porrati, Phys. Lett. B **498**, 92 (2001) [hep-th/0011152].

[50] A. I. Vainshtein, Phys. Lett. **39B**, 393 (1972).

[51] I. I. Kogan, S. Mouslopoulos, A. Papazoglou and G. G. Ross, Phys. Rev. D **64**, 124014 (2001) [hep-th/0107086].

[52] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Rev. D **63**, 075004 (2001) [hep-ph/0006041].

[53] S. Mouslopoulos and A. Papazoglou, JHEP **0011**, 018 (2000) [hep-ph/0003207].

[54] N. Kaloper, Phys. Lett. B **474**, 269 (2000) [hep-th/9912125].

[55] S. Nam, JHEP **0003**, 005 (2000) [hep-th/9911104].

[56] S. Nam, JHEP **0004**, 002 (2000) [hep-th/9911237].

[57] J. Lykken and L. Randall, JHEP **0006**, 014 (2000) [hep-th/9908076].

[58] P. Kanti, I. I. Kogan, K. A. Olive and M. Pospelov, Phys. Rev. D **61**, 106004 (2000) [hep-ph/9912266].

[59] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999) [hep-ph/9907447].

[60] Z. Chacko and P. J. Fox, hep-th/0102023.

[61] O. DeWolfe, D. Z. Freedman, S. S. Gubser and A. Karch, Phys. Rev. D **62**, 046008 (2000) [hep-th/9909134].

[62] P. Kanti, K. A. Olive and M. Pospelov, Phys. Lett. B **481**, 386 (2000) [hep-ph/0002229].

[63] N. Kaloper, Phys. Rev. D **60**, 123506 (1999) [hep-th/9905210].

[64] T. Nihei, Phys. Lett. B **465**, 81 (1999) [hep-ph/9905487].

[65] H. B. Kim and H. D. Kim, Phys. Rev. D **61**, 064003 (2000) [hep-th/9909053].

[66] P. Kanti, K. A. Olive and M. Pospelov, Phys. Rev. D **62**, 126004 (2000) [hep-ph/0005146].

[67] A. Miemiec, hep-th/0011160.

[68] E. D'Hoker, D. Z. Freedman, S. D. Mathur, A. Matusis and L. Rastelli, Nucl. Phys. B **562**, 330 (1999) [hep-th/9902042].

[69] A. Naqvi, JHEP **9912**, 025 (1999) [hep-th/9911182].

[70] I. L. Buchbinder, D. M. Gitman and V. D. Pershin, Phys. Lett. B **492**, 161 (2000) [hep-th/0006144].

[71] B. Allen and T. Jacobson, Commun. Math. Phys. **103**, 669 (1986).

[72] Y. M. Cho and S. W. Zoh, Phys. Rev. D **46**, 2290 (1992).

[73] T. Appelquist, A. Chodos and P. G. Freund, “Modern Kaluza-Klein Theories”, Addison-Wesley (1987).

[74] F. A. Dilkes, M. J. Duff, J. T. Liu and H. Sati, hep-th/0102093.

[75] M. J. Duff, J. T. Liu and H. Sati, hep-th/0105008.

[76] D. G. Boulware and S. Deser, Phys. Rev. D **6**, 3368 (1972).

[77] A. Karch, E. Katz and L. Randall, hep-th/0106261.

[78] C. Deffayet, G. Dvali, G. Gabadadze and A. Vainshtein, hep-th/0106001.

[79] A. Chodos and E. Poppitz, Phys. Lett. B **471**, 119 (1999) [hep-th/9909199].

[80] A. G. Cohen and D. B. Kaplan, Phys. Lett. B **470**, 52 (1999) [hep-th/9910132].

[81] R. Gregory, Phys. Rev. Lett. **84**, 2564 (2000) [hep-th/9911015].

[82] Z. Chacko and A. E. Nelson, Phys. Rev. D **62**, 085006 (2000) [hep-th/9912186].

[83] N. Arkani-Hamed, L. Hall, D. Smith and N. Weiner, Phys. Rev. D **62**, 105002 (2000) [hep-ph/9912453].

[84] J. Chen, M. A. Luty and E. Ponton, JHEP **0009**, 012 (2000) [hep-th/0003067].

[85] I. Olasagasti and A. Vilenkin, Phys. Rev. D **62**, 044014 (2000) [hep-th/0003300].

[86] T. Gherghetta and M. Shaposhnikov, Phys. Rev. Lett. **85**, 240 (2000) [hep-th/0004014].

[87] A. Chodos, E. Poppitz and D. Tsimpis, Class. Quant. Grav. **17**, 3865 (2000) [hep-th/0006093].

[88] T. Gherghetta, E. Roessl and M. Shaposhnikov, Phys. Lett. B **491**, 353 (2000) [hep-th/0006251].

[89] E. Ponton and E. Poppitz, JHEP **0102**, 042 (2001) [hep-th/0012033].

[90] S. Moon, S. Rey and Y. Kim, Nucl. Phys. B **602**, 467 (2001) [hep-th/0012165].

[91] C. Charmousis, R. Emparan and R. Gregory, JHEP **0105**, 026 (2001) [hep-th/0101198].

[92] O. Corradini and Z. Kakushadze, Phys. Lett. B **506**, 167 (2001) [hep-th/0103031].

[93] H. Collins and B. Holdom, hep-ph/0103103.

[94] J. E. Kim, B. Kyae and H. M. Lee, hep-th/0104150.

[95] P. Kanti, R. Madden and K. A. Olive, hep-th/0104177.

[96] S. Hayakawa and K. I. Izawa, hep-th/0106101.

[97] F. Leblond, R. C. Myers and D. J. Winters, hep-th/0106140.

[98] I. I. Kogan and N. A. Voronov, JETP Lett. **38**, 311 (1983).

[99] P. Candelas and S. Weinberg, Nucl. Phys. B **237**, 397 (1984).

[100] S. Deser, R. Jackiw and G. 't Hooft, Annals Phys. **152**, 220 (1984).

[101] S. Deser and R. Jackiw, Annals Phys. **153**, 405 (1984).

[102] I. I. Kogan, S. Mouslopoulos, A. Papazoglou and G. G. Ross, Nucl. Phys. B **615**, 191 (2001) [hep-ph/0107307].